

N Identical Particles Under Quantum Confinement: A Many-Body Dimensional Perturbation Theory Approach II, The Lowest-Order Wave Function I

M. Dunn, D. K. Watson

University of Oklahoma

Department of Physics and Astronomy

Norman, OK 73019

J. G. Loeser

Oregon State University

Department of Chemistry

Corvallis, OR 97331

February 1, 2008

Abstract

In this paper we continue our development of a dimensional perturbation theory (DPT) treatment of N identical particles under quantum confinement. DPT is a beyond-mean-field method which is applicable to both weakly and strongly-interacting systems and can be used to connect both limits. In a previous paper we developed the formalism for low-order energies and excitation frequencies. This formalism has been applied to atoms, Bose-Einstein condensates and quantum dots. One major advantage of the method is that N appears as a parameter in the analytical expressions for the energy and so results for N up to a few thousand are easy to obtain. Other properties however, are also of interest, for example the density profile in the case of a BEC, and larger N results are desirable as well. The latter case requires us to go to higher orders in DPT. These calculations require as input zeroth-order wave functions and this paper, along with a subsequent paper, address this issue.

1 Introduction

Systems that involve N identical interacting particles under quantum confinement appear throughout many areas of physics, including chemical, condensed matter, and atomic physics[1]. In a previous paper[2], we applied the methods of dimensional perturbation theory (DPT)[3], a powerful set of tools that uses symmetry to yield simple results, to such N -body systems. We presented a detailed discussion of the dimensional continuation of the N -particle Schrödinger equation, the $D \rightarrow \infty$ equilibrium (D^0) structure, and the (D^{-1}) energy. We used the FG matrix method to derive general, analytical expressions for the N -body normal-mode vibrational frequencies, and we gave specific analytical results for three confined N -body quantum

systems: the N -electron atom, the N -electron quantum dot, and the N -atom inhomogeneous Bose-Einstein condensate with a repulsive hardcore potential.

In a subsequent paper[4] we further pursued our analysis of the N -atom inhomogeneous Bose-Einstein condensate (BEC) with a repulsive hardcore potential and optimized our low-order DPT energy by fitting to low- N DMC data[5] and then extrapolated the result out to large N . Our low-order energies are accurate out to $N \approx 10,000$ and larger N could in principle be obtained by going to higher orders in the perturbation expansion[6]. Ten thousand atoms may not seem particularly impressive given the fact that the Gross-Pitaevskii (GP) equation can yield higher- N results, but the advantage of DPT is that it is not a weak-interatomic-interaction approximation. At lowest orders it includes beyond-mean-field effects. Thus DPT is effective for large- a systems that have been created in the lab by exploiting Feshbach resonances[7], systems for which the GP and modified Gross-Pitaevskii (MGP) equations fail. This we demonstrated for a BEC which had an a one hundred times that of the natural scattering length of ^{87}Rb .

In another paper[8], low-order N -body dimensional perturbation methods have been applied to the N -electron atom. In this instrumental paper, Loeser obtained low-order, analytical expressions for the ground-state energy of neutral atoms. For $Z = 1$ to 127, the numerical results compare well to Hartree-Fock energies with a correlation correction.

All of the previous work has focussed on low-order calculations of energies and thus far little attention has been given to wave functions, even at lowest order. Yet the lowest-order wave function yields important information about the system. The normal mode coordinates tell us the nature of the excitations of the system and the lowest-order wave function gives us expectation values at low orders. For macroscopic quantum-confined systems, such as the BEC, the wave function is uncloaked in an explicit way since the density profile may be viewed in a direct fashion. Also, calculating energies and wave functions to higher orders in $1/D$ requires as input the lowest-order wave function.

We briefly outlined the derivation of the lowest-order wave function in a previous letter[9] and in this paper we expand considerably upon that discussion. In Section 2 we discuss the S -wave, Jacobian-weighted Hamiltonian and wave function. In Section 3 we dimensionally scale the system to regularize the large dimension limit for which we then solve. In Section 4 we begin the normal mode analysis which will result in the energy through first order in $1/D$ and the lowest-order wave function. In Section 5 we discuss the symmetry of the N -body quantum-confinement problem. This symmetry simplifies the problem to an extraordinary degree, making an exact solution of this N -body problem, with $N(N - 1)/2$ interparticle interactions, possible at large D . In Section 6 we exploit this symmetry further and discuss the use of symmetry coordinates in the calculation of the large-dimension, zeroth-order normal modes of the system. In Section 7 we introduce a particular approach to determining a suitable set of symmetry coordinates. The method uses what we term primitive irreducible coordinates, from which the symmetry coordinates are derived. This is illustrated in detail in a simple example involving the coordinates of a particular species of the N -body quantum-confinement problem (symmetry coordinates transforming under the same irreducible

representation are said to belong to the same species). In Section 8 we apply the general theory discussed in earlier sections to derive in detail the frequencies and normal-mode coordinates of this species. Section 9 is a summary and conclusion section. The frequencies and normal-mode coordinates of the remaining, and more complicated species of the system are derived in a subsequent paper[10].

2 The D-dimensional N-body Schrödinger Equation

For an N -body system of particles confined by a spherically symmetric potential and interacting via a common two-body potential g_{ij} , the Schrödinger equation in D -dimensional Cartesian coordinates is

$$H\Psi = \left[\sum_{i=1}^N h_i + \sum_{i=1}^{N-1} \sum_{j=i+1}^N g_{ij} \right] \Psi = E\Psi, \quad (1)$$

where

$$h_i = -\frac{\hbar^2}{2m_i} \sum_{\nu=1}^D \frac{\partial^2}{\partial x_{i\nu}^2} + V_{\text{conf}} \left(\sqrt{\sum_{\nu=1}^D x_{i\nu}^2} \right), \quad (2)$$

$$\text{and } g_{ij} = V_{\text{int}} \left(\sqrt{\sum_{\nu=1}^D (x_{i\nu} - x_{j\nu})^2} \right) \quad (3)$$

are the single-particle Hamiltonian and the two-body interaction potential, respectively. The operator H is the D -dimensional Hamiltonian, and $x_{i\nu}$ is the ν^{th} Cartesian component of the i^{th} particle. The term V_{conf} is the confining potential. For the N -electron atom, V_{conf} is the Coulomb attraction between the electrons and the nucleus, while for the N -electron quantum dot and N -atom hard-sphere problem, a model appropriate for trapped gaseous BECs, we model the confinement as a harmonic trapping potential. The two-body interaction potential V_{int} is Coulombic in the first two systems and a hard sphere in the third.

2.1 Transformation of the Laplacian

Restricting our attention to spherically symmetric ($L = 0$) states of the N -body system, we transform from Cartesian to internal coordinates. A convenient internal coordinate system for confined systems is

$$r_i = \sqrt{\sum_{\nu=1}^D x_{i\nu}^2} \quad (1 \leq i \leq N) \quad \text{and} \quad \gamma_{ij} = \cos(\theta_{ij}) = \left(\sum_{\nu=1}^D x_{i\nu} x_{j\nu} \right) / r_i r_j \quad (1 \leq i < j \leq N), \quad (4)$$

which are the D -dimensional scalar radii r_i of the N particles from the center of the confining potential and the cosines γ_{ij} of the $N(N-1)/2$ angles between the radial vectors.

Now for a function Ψ dependent on only two functions $r(x)$ and $\gamma(x)$ one can write

$$\frac{d^2 \Psi(r(x), \gamma(x))}{dx^2} = \frac{d^2 r}{dx^2} \frac{d\Psi}{dr} + \frac{d^2 \gamma}{dx^2} \frac{d\Psi}{d\gamma} + \left(\frac{dr}{dx} \right)^2 \frac{d^2 \Psi}{dr^2} + \left(\frac{d\gamma}{dx} \right)^2 \frac{d^2 \Psi}{d\gamma^2} + 2 \frac{d\gamma}{dx} \frac{dr}{dx} \frac{d^2 \Psi}{dr d\gamma}. \quad (5)$$

Generalizing this, when operating on the state $\Psi(r_i(x_{i\nu}), \gamma_{i1}(x_{i\nu}) \dots \gamma_{ik}(x_{i\nu}) \dots \gamma_{iN}(x_{i\nu}))$ (where $k \neq i$ and $\nu = 1, \dots, D$), ∇_i^2 can be written in terms of the internal coordinates of Eq. (4) as

$$\begin{aligned} \nabla_i^2 \Psi \equiv \sum_{\nu=1}^D \frac{\partial^2}{\partial x_{i\nu}^2} \Psi &= \sum_{\nu=1}^D \left(\frac{\partial^2 r_i}{\partial x_{i\nu}^2} \right) \frac{\partial}{\partial r_i} \Psi + \sum_{\nu=1}^D \sum_{j \neq i} \left(\frac{\partial^2 \gamma_{ij}}{\partial x_{i\nu}^2} \right) \frac{\partial}{\partial \gamma_{ij}} \Psi + \\ &\sum_{\nu=1}^D \left(\frac{\partial r_i}{\partial x_{i\nu}} \right)^2 \frac{\partial^2}{\partial r_i^2} \Psi + \sum_{\nu=1}^D \sum_{j \neq i} \sum_{k \neq i} \left(\frac{\partial \gamma_{ij}}{\partial x_{i\nu}} \right) \left(\frac{\partial \gamma_{ik}}{\partial x_{i\nu}} \right) \frac{\partial^2}{\partial \gamma_{ij} \partial \gamma_{ik}} \Psi + \\ &2 \sum_{\nu=1}^D \sum_{j \neq i} \left(\frac{\partial r_i}{\partial x_{i\nu}} \right) \left(\frac{\partial \gamma_{ij}}{\partial x_{i\nu}} \right) \frac{\partial^2}{\partial r_i \partial \gamma_{ij}} \Psi. \end{aligned} \quad (6)$$

The relevant derivatives of the internal coordinates are

$$\frac{\partial r_i}{\partial x_{i\nu}} = \frac{x_{i\nu}}{r_i} \quad \frac{\partial \gamma_{ij}}{\partial x_{i\nu}} = \frac{1}{r_i} \left(\frac{x_{j\nu}}{r_j} - \frac{x_{i\nu}}{r_i} \gamma_{ij} \right) \quad (7)$$

$$\frac{\partial^2 r_i}{\partial x_{i\nu}^2} = \frac{1}{r_i} \left(1 - \frac{x_{i\nu}^2}{r_i^2} \right) \quad \frac{\partial^2 \gamma_{ij}}{\partial x_{i\nu}^2} = \frac{1}{r_i^2} \left(3 \frac{x_{i\nu}^2}{r_i^2} \gamma_{ij} - 2 \frac{x_{i\nu}}{r_i} \frac{x_{j\nu}}{r_j} - \gamma_{ij} \right), \quad (8)$$

which lead to the effective S -wave Laplacian in internal coordinates:

$$\begin{aligned} \sum_i \nabla_i^2 \Psi &= \sum_i \frac{D-1}{r_i} \frac{\partial}{\partial r_i} \Psi - \sum_i \frac{D-1}{r_i^2} \sum_{j \neq i} \gamma_{ij} \frac{\partial}{\partial \gamma_{ij}} \Psi + \\ &\sum_i \frac{\partial^2}{\partial r_i^2} \Psi + \sum_i \sum_{j \neq i} \sum_{k \neq i} \frac{\gamma_{jk} - \gamma_{ij} \gamma_{ik}}{r_i^2} \frac{\partial^2}{\partial \gamma_{ij} \partial \gamma_{ik}} \Psi. \end{aligned} \quad (9)$$

2.2 The Jacobian-Weighted Hamiltonian

All of the N -body DPT work up till now has utilized a similarity transformation so that the kinetic energy operator is transformed to a sum of terms of two kinds, namely, derivative terms *and* a repulsive centrifugal-like term, which when attractive interparticle potentials are present, stabilizes the system against collapse in the large- D limit. The particular transformation which has been used in almost all the work to date, is chosen so that first-order derivatives resulting from the Laplacian (Eq. (9)) are removed from the transformed Hamiltonian. When this is done the zeroth and first orders of the dimensional ($1/D$) expansion of the Hamiltonian become exactly soluble for any value of N . In the $D \rightarrow \infty$ limit, the second derivative terms drop out, resulting in a static problem at zeroth order, while first order corrections correspond to simple harmonic normal-mode oscillations about the infinite-dimensional structure.

Writing the similarity transformation of the wave function Ψ and operators $\hat{O}$ as

$$\Phi = \chi^{-1} \Psi, \quad \text{and} \quad \tilde{O} = \chi^{-1} \hat{O} \chi, \quad (10)$$

the transforming function is chosen to be

$$\chi = (r_1 r_2 \dots r_N)^{-(D-1)/2} \Gamma^{-(D-1)/4}. \quad (11)$$

where Γ is the Gramian determinant, the determinant of the matrix whose elements are γ_{ij} (see Appendix B).

Carrying out the transformation of the Hamiltonian and wave function of Eq. (1) via Eqs. (10) and (11), the Schrödinger equation becomes[8]:

$$({}_{(iii)}T + V) {}_{(iii)}\Phi = E {}_{(iii)}\Phi \quad (12)$$

$$({}_{(iii)}T = \sum_{i=1}^N \left(-\frac{\hbar^2}{2m_i} \frac{\partial^2}{\partial r_i^2} - \sum_{j \neq i} \sum_{k \neq i} \frac{\hbar^2(\gamma_{jk} - \gamma_{ij}\gamma_{ik})}{2m_i r_i^2} \frac{\partial^2}{\partial \gamma_{ij} \partial \gamma_{ik}} + \frac{\hbar^2(D-1)(D-2N-1)}{8m_i r_i^2} \frac{\Gamma^{(i)}}{\Gamma} \right) \quad (13)$$

$$V = \sum_{i=1}^N V_{\text{conf}}(r_i) + \sum_{i=1}^{N-1} \sum_{j=i+1}^N V_{\text{int}}(r_{ij}). \quad (14)$$

The Gramian matrix whose determinant is $\Gamma^{(i)}$ is the i^{th} principal minor formed by deleting from Γ the row and column corresponding to the i^{th} particle. The quantity $r_{ij} = \sqrt{r_i^2 + r_j^2 - 2r_i r_j \gamma_{ij}}$ is the interparticle separation. The similarity transformed Hamiltonian for the energy eigenstate ${}_{(iii)}\Phi$ is ${}_{(iii)}H = \chi^{-1} H \chi$, where ${}_{(iii)}H = ({}_{(iii)}T + V)$. As we can see from Eq. (13) all first-order derivatives have been eliminated from the Hamiltonian.

While the similarity transformation of Eqs. (10) and (11) is convenient in that all first-order derivative terms are eliminated, it complicates the interpretation of the normal-mode structure of the large- D wave function. Only when the weight function, W , for the matrix elements is equal to unity, i.e.

$$W = J \chi^2 = 1 \quad (15)$$

or

$$\chi = J^{-1/2}, \quad (16)$$

where J is the Jacobian of the transformation to internal coordinates, will a first derivative of an internal coordinate be the conjugate momentum to that coordinate. The transforming function of Eq. (11) does not satisfy Eq. (16) since the Jacobian J is[11]

$$J = (r_1 r_2 \dots r_N)^{D-1} \Gamma^{(D-N-1)/2}. \quad (17)$$

Furthermore, the development of higher-order DPT expansions involves the matrix elements of coordinates and their derivatives between the zeroth-order normal-mode functions. These matrix elements are much easier to calculate when the weight function in the integrals is unity, and this only occurs when Eq (16) is satisfied.

In Ref. [11], Avery *et al.* considered the problem of performing the similarity transformation of Eq. (10) with a more general transforming function χ , which has adjustable parameters α and β , and is of the form:

$$\chi = (r_1 r_2 \dots r_N)^{-\alpha} \Gamma^{-\beta/2}. \quad (18)$$

One of the cases considered by Avery *et al.*[11] has $\alpha = (D-1)/2$ and $\beta = (D-N-1)/2$ so that

$$\chi = (r_1 r_2 \dots r_N)^{-(D-1)/2} \Gamma^{-(D-N-1)/4}. \quad (19)$$

The transforming function χ of Eq. (19) does satisfy Eqs. (15) and (16).

Carrying out the transformation of the Hamiltonian and wave function of Eq. (1) via Eqs. (10) and (19), the Schrödinger equation becomes[11]:

$$({}_i T + V) {}_i \Phi = E {}_i \Phi \quad (20)$$

where

$$\begin{aligned} {}_i T &= \hbar^2 \sum_{i=1}^N \left[-\frac{1}{2m_i} \frac{\partial^2}{\partial r_i^2} - \frac{1}{2m_i r_i^2} \left(\sum_{j \neq i} \sum_{k \neq i} (\gamma_{jk} - \gamma_{ij} \gamma_{ik}) \frac{\partial^2}{\partial \gamma_{ij} \partial \gamma_{ik}} - N \sum_{j \neq i} \gamma_{ij} \frac{\partial}{\partial \gamma_{ij}} \right) \right. \\ &\quad \left. + \frac{N(N-2) + (D-N-1)^2 \left(\frac{\Gamma^{(i)}}{\Gamma} \right)}{8m_i r_i^2} \right] \\ &= \hbar^2 \sum_{i=1}^N \left[-\frac{1}{2m_i} \frac{\partial^2}{\partial r_i^2} - \frac{1}{2m_i r_i^2} \sum_{j \neq i} \sum_{k \neq i} \frac{\partial}{\partial \gamma_{ij}} (\gamma_{jk} - \gamma_{ij} \gamma_{ik}) \frac{\partial}{\partial \gamma_{ik}} \right. \\ &\quad \left. + \frac{N(N-2) + (D-N-1)^2 \left(\frac{\Gamma^{(i)}}{\Gamma} \right)}{8m_i r_i^2} \right] \end{aligned} \quad (21)$$

The latter expression for ${}_i T$ is explicitly self-adjoint since the weight function, W , for the matrix elements is equal to unity, i.e. Eq. (15) is satisfied. The similarity transformed Hamiltonian for the energy eigenstate ${}_i \Phi$ is ${}_i H = ({}_i T + V)$.

3 Infinite-D analysis: Leading order energy

To begin the perturbation analysis we regularize the large-dimension limit of the Schrödinger equation by defining dimensionally scaled variables:

$$\bar{r}_i = r_i / \kappa(D) \quad , \quad \bar{E} = \kappa(D) E \quad \text{and} \quad \bar{H} = \kappa(D) {}_i H \quad (22)$$

with dimension-dependent scale factor $\kappa(D)$. From Eq. (21) one can see that the kinetic energy T scales in the same way as $1/r^2$, so the scaled version of Eq. (20) becomes

$$\bar{H} \Phi = \left(\frac{1}{\kappa(D)} \bar{T} + \bar{V}_{\text{eff}} \right) \Phi = \bar{E} \Phi, \quad (23)$$

where barred quantities simply indicate that the variables are now in scaled units. Because of the quadratic D dependence in the centrifugal-like term in T of Eq. (21), we conclude that the scale factor $\kappa(D)$ must also be quadratic in D , otherwise the $D \rightarrow \infty$ limit of the Hamiltonian would not be finite. The precise form of $\kappa(D)$ is chosen so that the result of the scaling is as simple as possible and depends on the system in question. In previous work[2] we have chosen $\kappa(D) = (D-1)(D-2N-1)/(4Z)$ for the S -wave, N -electron atom; $\kappa(D) = \Omega l_{\text{ho}}$ for the N -electron quantum dot where $\Omega = (D-1)(D-2N-1)/4$ and the dimensionally-scaled harmonic oscillator length and trap frequency respectively are $l_{\text{ho}} = \sqrt{\frac{\hbar}{m^* \bar{\omega}_{\text{ho}}}}$ and $\bar{\omega}_{\text{ho}}^2 = \Omega^3 \omega_{\text{ho}}^2$; and $\kappa(D) = D^2 \bar{a}_{\text{ho}}$ for the BEC where $\bar{a}_{\text{ho}} = \sqrt{\frac{\hbar}{m \bar{\omega}_{\text{ho}}}}$ and $\bar{\omega}_{\text{ho}} = D^3 \omega_{\text{ho}}$. The factor of $\kappa(D)$ in the denominator of

Eq. (23) acts as an effective mass that increases with D , causing the derivative terms to become suppressed while leaving behind a centrifugal-like term in an effective potential,

$$\bar{V}_{\text{eff}}(\bar{r}, \gamma; \delta = 0) = \sum_{i=1}^N \left(\frac{\hbar^2}{8m_i \bar{r}_i^2} \frac{\Gamma^{(i)}}{\Gamma} + \bar{V}_{\text{conf}}(\bar{r}, \gamma; \delta = 0) \right) + \sum_{i=1}^{N-1} \sum_{j=i+1}^N \bar{V}_{\text{int}}(\bar{r}, \gamma; \delta = 0), \quad (24)$$

where $\delta = 1/D$, in which the particles become frozen at large D . In the $D \rightarrow \infty$ ($\delta \rightarrow 0$) limit, the excited states have collapsed onto the ground state, which is found at the minimum of V_{eff} .

We assume a totally symmetric minimum characterized by the equality of all radii and angle cosines of the particles when $D \rightarrow \infty$, i.e.

$$\bar{r}_i = \bar{r}_\infty \quad (1 \leq i \leq N), \quad \gamma_{ij} = \bar{\gamma}_\infty \quad (1 \leq i < j \leq N). \quad (25)$$

Since each particle radius and angle cosine is equivalent, we can take derivatives with respect to an arbitrary $\bar{r}_i$ and γ_{ij} in the minimization procedure. Then evaluating all $\bar{r}_i$ and γ_{ij} at the infinite- D radius and angle cosine, $\bar{r}_\infty$ and $\bar{\gamma}_\infty$, respectively, we find that $\bar{r}_\infty$ and $\bar{\gamma}_\infty$ satisfy

$$\left[\frac{\partial \bar{V}_{\text{eff}}(\bar{r}, \gamma; \delta)}{\partial \bar{r}_i} \right]_{\delta=0} = 0 \quad (26)$$

$$\left[\frac{\partial \bar{V}_{\text{eff}}(\bar{r}, \gamma; \delta)}{\partial \gamma_{ij}} \right]_{\delta=0} = 0, \quad (27)$$

where the $\delta = 0$ subscript means to evaluate all $\bar{r}_i$ at $\bar{r}_\infty$ and all γ_{ij} at $\bar{\gamma}_\infty$. In scaled units the zeroth-order ($D \rightarrow \infty$) approximation for the energy becomes

$$\bar{E}_\infty = \bar{V}_{\text{eff}}(\bar{r}_\infty, \bar{\gamma}_\infty; 0). \quad (28)$$

In this leading order approximation, the centrifugal-like term that appears in $\bar{V}_{\text{eff}}$, even for the ground state, is a zero-point energy contribution required by the minimum uncertainty principle[12].

4 Normal-mode analysis and the 1/D first-order quantum energy correction

At zeroth-order, the particles can be viewed as frozen in a completely symmetric, high- D configuration or simplex, which is somewhat analogous to the Lewis structure in atomic physics terminology. Likewise, the first-order $1/D$ correction can be viewed as small oscillations of this structure, analogous to Langmuir oscillations. Solving Eqs. (26) and (27) for $\bar{r}_\infty$ and $\bar{\gamma}_\infty$ gives the infinite- D structure and zeroth-order energy and provides the starting point for the $1/D$ expansion. To obtain the $1/D$ quantum correction to the energy for large but finite values of D , we expand about the minimum of the $D \rightarrow \infty$ effective potential. We first define a position vector, consisting of all $N(N+1)/2$ internal coordinates:

$$\bar{\mathbf{y}} = \begin{pmatrix} \bar{\mathbf{r}} \\ \gamma \end{pmatrix}, \quad (29)$$

where

$$\bar{\mathbf{r}} = \begin{pmatrix} \bar{r}_1 \\ \bar{r}_2 \\ \vdots \\ \bar{r}_N \end{pmatrix} \quad (30)$$

and

$$\boldsymbol{\gamma} = \begin{pmatrix} \gamma_{12} \\ \gamma_{13} \\ \gamma_{23} \\ \gamma_{14} \\ \gamma_{24} \\ \gamma_{34} \\ \gamma_{15} \\ \gamma_{25} \\ \vdots \\ \gamma_{N-2,N} \\ \gamma_{N-1,N} \end{pmatrix}. \quad (31)$$

We then make the following substitutions for all radii and angle cosines:

$$\bar{r}_i = \bar{r}_\infty + \delta^{1/2} \bar{r}'_i \quad (32)$$

$$\gamma_{ij} = \bar{\gamma}_\infty + \delta^{1/2} \bar{\gamma}'_{ij}, \quad (33)$$

where $\delta = 1/D$ is the expansion parameter. Now in all practical situations $\bar{V}_{\text{eff}}(\bar{y}; \delta)$ is a function of $\bar{y}$ and δ and so we may then obtain a power series in $\delta^{1/2}$ of the effective potential about the $D \rightarrow \infty$ symmetric minimum as

$$\begin{aligned} \bar{V}_{\text{eff}}(\bar{y}; \delta) &= [\bar{V}_{\text{eff}}]_{\delta^{1/2}=0} + \\ &+ \delta^{1/2} \left\{ \left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \delta} \right]_{\bar{y}_\mu} \Big|_{\delta^{1/2}=0} \left[\frac{\partial \delta}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} + \sum_{\mu=1}^P \left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu} \right]_{\delta^{1/2}=0} \left[\frac{\partial \bar{y}_\mu}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} \right\} + \\ &+ \frac{1}{2} \delta \left\{ \left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \delta} \right]_{\bar{y}_\mu} \Big|_{\delta^{1/2}=0} \left[\frac{\partial^2 \delta}{(\partial \delta^{1/2})^2} \right]_{\delta^{1/2}=0} + \left[\frac{\partial^2 \bar{V}_{\text{eff}}}{\partial \delta^2} \right]_{\bar{y}_\mu} \Big|_{\delta^{1/2}=0} \left[\frac{\partial \delta}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0}^2 + \right. \\ &+ \sum_{\mu=1}^P \left(2 \left[\frac{\partial}{\partial \delta} \right]_{\bar{y}_\mu} \left(\left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu} \right]_{\delta^{1/2}} \right) \right]_{\delta^{1/2}=0} \left[\frac{\partial \bar{y}_\mu}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} \left[\frac{\partial \delta}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} + \\ &+ \left. \left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu} \right]_{\delta^{1/2}} \Big|_{\delta^{1/2}=0} \left[\frac{\partial^2 \bar{y}_\mu}{(\partial \delta^{1/2})^2} \right]_{\delta^{1/2}=0} \right) + \\ &+ \sum_{\mu=1}^P \sum_{\nu=1}^P \left[\frac{\partial \bar{y}_\mu}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} \left[\frac{\partial^2 \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu \partial \bar{y}_\nu} \right]_{\delta^{1/2}=0} \left[\frac{\partial \bar{y}_\nu}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} \Big\} + O(\delta^{3/2}), \end{aligned} \quad (34)$$

where

$$P \equiv N(N+1)/2 \quad (35)$$

is the number of internal coordinates. The $O((\delta^{1/2})^0)$ term in the power series (Eq. 43) is simply the zeroth-order energy (Eq. 28). The $O((\delta^{1/2})^1)$ term is zero since

$$\left[\frac{\partial \delta}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} = 0 \quad (36)$$

and we are expanding about the minimum of the effective potential so

$$\left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu} \right]_{\delta^{1/2}=0} = 0, \quad (37)$$

i.e. the system is said to be in equilibrium since the forces acting on the system vanish [Eqs. (26) and (27)].

Defining a displacement vector consisting of the internal displacement coordinates [primed in Eqs. (32) and (33)]

$$\bar{\mathbf{y}}' = \begin{pmatrix} \bar{\mathbf{r}}' \\ \bar{\boldsymbol{\gamma}}' \end{pmatrix}, \quad (38)$$

where

$$\bar{\mathbf{r}}' = \begin{pmatrix} \bar{r}'_1 \\ \bar{r}'_2 \\ \vdots \\ \bar{r}'_N \end{pmatrix} \quad (39)$$

and

$$\bar{\boldsymbol{\gamma}}' = \begin{pmatrix} \bar{\gamma}'_{12} \\ \bar{\gamma}'_{13} \\ \bar{\gamma}'_{23} \\ \bar{\gamma}'_{14} \\ \bar{\gamma}'_{24} \\ \bar{\gamma}'_{34} \\ \bar{\gamma}'_{15} \\ \bar{\gamma}'_{25} \\ \vdots \\ \bar{\gamma}'_{N-2,N} \\ \bar{\gamma}'_{N-1,N} \end{pmatrix}, \quad (40)$$

we then have

$$\left[\frac{\partial \bar{y}_\mu}{\partial \delta^{1/2}} \right]_{\delta^{1/2}=0} = \bar{y}'_\mu \quad (41)$$

and also defining

$$v_o = \left[\frac{\partial \bar{V}_{\text{eff}}}{\partial \delta} \right]_{\bar{y}_\mu}_{\delta^{1/2}=0}, \quad (42)$$

Eq. (34) becomes

$$\bar{V}_{\text{eff}}(\bar{\mathbf{y}}'; \delta) = [\bar{V}_{\text{eff}}]_{\delta^{1/2}=0} + \frac{1}{2} \delta \left\{ \sum_{\mu=1}^P \sum_{\nu=1}^P \bar{y}'_\mu \left[\frac{\partial^2 \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu \partial \bar{y}_\nu} \right]_{\delta^{1/2}=0} \bar{y}'_\nu + v_o \right\} + O(\delta^{3/2}), \quad (43)$$

The first term of the $O((\delta^{1/2})^2)$ term defines the elements of the Hessian matrix[13] $\mathbf{F}$ of Eq. (45) below. The derivative terms in the kinetic energy are taken into account by a similar series expansion, beginning with a first-order term that is bilinear in $\partial/\partial\bar{y}'$, i.e.

$$\mathcal{T} = -\frac{1}{2}\delta \sum_{\mu=1}^P \sum_{\nu=1}^P G_{\mu\nu} \partial_{\bar{y}'_\mu} \partial_{\bar{y}'_\nu} + O\left(\delta^{3/2}\right), \quad (44)$$

where $\mathcal{T}$ is the derivative portion of the kinetic energy T (see Eq. (21)). Thus, obtaining the first-order energy correction is reduced to a harmonic problem, which is solved by obtaining the normal modes of the system.

We use the Wilson FG matrix method[14] to obtain the normal-mode vibrations and, thereby, the first-order energy correction. It follows from Eqs. (43) and (44) that $\mathbf{G}$ and $\mathbf{F}$, both constant matrices, are defined in the first-order $\delta = 1/D$ Hamiltonian as follows:

$$\hat{H}_1 = -\frac{1}{2}\partial_{\bar{y}'}^T \mathbf{G} \partial_{\bar{y}'} + \frac{1}{2}\bar{\mathbf{y}}'^T \mathbf{F} \bar{\mathbf{y}}' + v_o. \quad (45)$$

After the Schrödinger equation (20) has been dimensionally scaled, the second-order derivative terms are of order δ , and by comparing these terms with the first part of $\hat{H}_1$, the elements of the kinetic-energy matrix $\mathbf{G}$ are easily determined. The elements of the Hessian matrix[13], $F_{\mu\nu} = \left[\frac{\partial^2 \bar{V}_{\text{eff}}}{\partial \bar{y}_\mu \partial \bar{y}_\nu} \right]_{\delta^{1/2}=0}$, on the other hand, require a bit more effort to obtain[2].

We include a derivation of the FG matrix method in Appendix A, but we state here the main results of the method. The b^{th} normal mode coordinate may be written as (Eq. (A19))

$$[\mathbf{q}']_b = \mathbf{b}^T \bar{\mathbf{y}}', \quad (46)$$

where the coefficient vector $\mathbf{b}$ satisfies the eigenvalue equation (Eq. (A20))

$$\mathbf{F} \mathbf{G} \mathbf{b} = \lambda_b \mathbf{b} \quad (47)$$

with the resultant secular equation (Eq. (A21))

$$\det(\mathbf{F} \mathbf{G} - \lambda \mathbf{I}) = 0. \quad (48)$$

The coefficient vector also satisfies the normalization condition (Eq. (A22))

$$\mathbf{b}^T \mathbf{G} \mathbf{b} = 1. \quad (49)$$

As can be seen from Eq. (A5) in Appendix A the frequencies are given by

$$\lambda_b = \bar{\omega}_b^2, \quad (50)$$

while the wave function is a product of $P = N(N+1)/2$ harmonic oscillator wave functions

$$\Phi_0(\bar{\mathbf{y}}') = \prod_{b=1}^P h_{n_b} \left(\bar{\omega}_b^{1/2} [\mathbf{q}']_b \right), \quad (51)$$

where $h_{n_b}(\bar{\omega}_b^{1/2}[\mathbf{q}']_b)$ is a one-dimensional harmonic-oscillator wave function of frequency $\bar{\omega}_b$, and n_b is the oscillator quantum number, $0 \leq n_b < \infty$, which counts the number of quanta in each normal mode.

Depending on the number of particles, the number of roots λ of Eq. (48) – there are $P \equiv N(N+1)/2$ roots – is potentially huge. However, due to the S_N symmetry (see Ref. [15] and App. C) of the problem (see Sect. 5), there is a remarkable simplification. Equation (48) has only five distinct roots, λ_μ , where μ runs over $\mathbf{0}^-$, $\mathbf{0}^+$, $\mathbf{1}^-$, $\mathbf{1}^+$, and $\mathbf{2}$, regardless of the number of particles in the system (see Refs. [2, 8] and Sect. 6.3). Thus the energy through first-order (see Eq. (53)) can be written in terms of the five distinct normal-mode vibrational frequencies, which are related to the roots λ_μ of $\mathbf{FG}$ by

$$\lambda_\mu = \bar{\omega}_\mu^2. \quad (52)$$

The energy through first-order in $\delta = 1/D$ is then[2]

$$\bar{E} = \bar{E}_\infty + \delta \left[\sum_{\mu=\{\mathbf{0}^\pm, \mathbf{1}^\pm, \mathbf{2}\}} \sum_{n_\mu=0}^{\infty} (n_\mu + \frac{1}{2}) d_{\mu, n_\mu} \bar{\omega}_\mu + v_o \right], \quad (53)$$

where the n_μ are the vibrational quantum numbers of the normal modes of the same frequency $\bar{\omega}_\mu$ (as such, n_μ counts the number of nodes in a given normal mode). The quantity d_{μ, n_μ} is the occupancy of the manifold of normal modes with vibrational quantum number n_μ and normal mode frequency $\bar{\omega}_\mu$, i.e. it is the number of normal modes with the same frequency $\bar{\omega}_\mu$ and the same number of quanta n_μ . The total occupancy of the normal modes with frequency $\bar{\omega}_\mu$ is equal to the multiplicity of the root λ_μ , i.e.

$$d_\mu = \sum_{n_\mu=0}^{\infty} d_{\mu, n_\mu}, \quad (54)$$

where d_μ is the multiplicity of the μ^{th} root. The multiplicities of the five roots are[2]

$$\begin{aligned} d_{\mathbf{0}^+} &= 1, \\ d_{\mathbf{0}^-} &= 1, \\ d_{\mathbf{1}^+} &= N-1, \\ d_{\mathbf{1}^-} &= N-1, \\ d_{\mathbf{2}} &= N(N-3)/2. \end{aligned} \quad (55)$$

Note that although the equation in Ref. [8] for the energy through Langmuir order is the same as Eq. (53), it is expressed a little differently (See Ref. [16]).

5 The Symmetry of the Large-D, N-body Quantum-Confinement Problem

Such a high degree of degeneracy of the large- D , N -body quantum-confinement problem indicates a high degree of symmetry. In this section we study this in more detail.

5.1 The Indical structure of $\mathbf{F}$, $\mathbf{G}$, and $\mathbf{FG}$ matrices

The $\mathbf{F}$, $\mathbf{G}$, and $\mathbf{FG}$ matrices, which we generically denote by $\mathbf{Q}$, are $P \times P$ matrices with the same indicial structure as $\bar{y}\bar{y}^T$:

$$\bar{y}\bar{y}^T = \left(\begin{array}{cccc|cccc} \bar{r}'_1 \bar{r}'_1 & \bar{r}'_1 \bar{r}'_2 & \cdots & \bar{r}'_1 \bar{r}'_N & \bar{r}'_1 \bar{\gamma}'_{12} & \bar{r}'_1 \bar{\gamma}'_{13} & \cdots & \bar{r}'_1 \bar{\gamma}'_{N-1N} \\ \bar{r}'_2 \bar{r}'_1 & \bar{r}'_2 \bar{r}'_2 & \cdots & & \bar{r}'_2 \bar{\gamma}'_{12} & \bar{r}'_2 \bar{\gamma}'_{13} & \cdots & \bar{r}'_2 \bar{\gamma}'_{N-1N} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \bar{r}'_N \bar{r}'_1 & \cdots & & \bar{r}'_N \bar{r}'_N & \bar{r}'_N \bar{\gamma}'_{12} & & \cdots & \bar{r}'_N \bar{\gamma}'_{N-1N} \\ \hline \bar{\gamma}'_{12} \bar{r}'_1 & \bar{\gamma}'_{12} \bar{r}'_2 & \cdots & \bar{\gamma}'_{12} \bar{r}'_N & \bar{\gamma}'_{12} \bar{\gamma}'_{12} & \bar{\gamma}'_{12} \bar{\gamma}'_{13} & \cdots & \bar{\gamma}'_{12} \bar{\gamma}'_{N-1N} \\ \bar{\gamma}'_{13} \bar{r}'_1 & \bar{\gamma}'_{13} \bar{r}'_2 & \cdots & \bar{\gamma}'_{13} \bar{r}'_N & \bar{\gamma}'_{13} \bar{\gamma}'_{12} & \bar{\gamma}'_{13} \bar{\gamma}'_{13} & \cdots & \bar{\gamma}'_{13} \bar{\gamma}'_{N-1N} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \bar{\gamma}'_{N-1N} \bar{r}'_1 & \cdots & & \bar{\gamma}'_{N-1N} \bar{r}'_N & \bar{\gamma}'_{N-1N} \bar{\gamma}'_{12} & \bar{\gamma}'_{N-1N} \bar{\gamma}'_{13} & \cdots & \bar{\gamma}'_{N-1N} \bar{\gamma}'_{N-1N} \end{array} \right), \quad (56)$$

where $\bar{y}$ is defined by Eq. (29). The indicial structure of this matrix suggests a convenient shorthand for referencing the elements of the $\mathbf{Q}$ matrices. The upper left block of Eq. (56) is an $(N \times N)$ matrix with elements associated with $(\bar{r}'_i, \bar{r}'_j)$; hence we use the subscript (i, j) to refer to these elements. The upper right block is an $(N \times N(N-1)/2)$ matrix with elements associated with $(\bar{r}'_i, \bar{\gamma}'_{jk})$; hence, we use the subscript (i, jk) to refer to these elements. The lower left block is an $(N(N-1)/2 \times N)$ matrix with elements associated with $(\bar{\gamma}'_{ij}, \bar{r}'_k)$; hence, we use the subscript (ij, k) to refer to these elements. Finally, the lower right block is an $(N(N-1)/2 \times N(N-1)/2)$ matrix with elements associated with $(\bar{\gamma}'_{ij}, \bar{\gamma}'_{kl})$; hence, we use the subscript (ij, kl) to refer to these elements.

5.2 The S_N symmetry of the $\mathbf{Q}$ matrices

As the number of particles N increases, diagonalizing the $P \times P$ $\mathbf{FG}$ matrix (where $P \equiv N(N+1)/2$) becomes, prima facie, a daunting task. However, one of the advantages of dimensional perturbation theory is the simplifications that occur in the large-dimension limit. In particular, since we are dealing with identical particles in a totally symmetric configuration (the Lewis structure) in which all the particles are equivalent, the $\mathbf{Q}$ matrices display a high degree of symmetry with many identical elements. Specifically,

$$\begin{aligned} Q_{i,i} &= Q_{i',i'} &\equiv Q_a \\ Q_{i,j} &= Q_{i',j'} &\equiv Q_b \quad (i \neq j) \text{ and } (i' \neq j') \\ Q_{ij,i} &= Q_{i'j',i'} &\equiv Q_c \quad (i \neq j) \text{ and } (i' \neq j') \\ Q_{jk,i} &= Q_{j'k',i'} &\equiv Q_d \quad (i \neq j \neq k) \text{ and } (i' \neq j' \neq k') \\ Q_{i,ij} &= Q_{i',i'j'} &\equiv Q_e \quad (i \neq j) \text{ and } (i' \neq j') \\ Q_{i,jk} &= Q_{i',j'k'} &\equiv Q_f \quad (i \neq j \neq k) \text{ and } (i' \neq j' \neq k') \\ Q_{ij,ij} &= Q_{i'j',i'j'} &\equiv Q_g \quad (i \neq j) \text{ and } (i' \neq j') \\ Q_{ij,jk} &= Q_{i'j',j'k'} &\equiv Q_h \quad (i \neq j \neq k) \text{ and } (i' \neq j' \neq k') \\ Q_{ij,kl} &= Q_{i'j',k'l'} &\equiv Q_i \quad (i \neq j \neq k \neq l) \text{ and } (i' \neq j' \neq k' \neq l'). \end{aligned} \quad (57)$$

Note the indices in the relationships above run over all particles $(1, 2, \dots, N)$ with the exceptions noted in the far right column. For example, $Q_{i,j} = Q_{i',j'} \equiv Q_b$, where $(i \neq j)$ and $(i' \neq j')$, means that all off-diagonal elements of the upper left block (the pure radial block) of $\mathbf{Q}$ are equal to the same constant Q_b . Similarly, $Q_{ij,kl} = Q_{i'j',k'l'} \equiv Q_\iota$, where $(i \neq j \neq k \neq l)$ and $(i' \neq j' \neq k' \neq l')$, means that any elements of $\mathbf{Q}$ in the lower right block (the pure angular block) that do not have a repeated index are all equal to the same constant Q_ι .

Equations (57) imply that the $\mathbf{Q}$ matrices, $\mathbf{F}$, $\mathbf{G}$, and $\mathbf{FG}$, are invariant under interchange of the particle labels. Thus Eqs. (57) show that the system is invariant under the group of all the permutations of N objects, where the objects are the particle label indices. This is the symmetric group S_N (see Appendix C). As we shall see, this S_N invariance greatly simplifies the problem which would otherwise be intractable for large N ; allowing us to solve for the $N(N+1)/2$ normal-mode coordinates, and hence the Langmuir-order wave function. The S_N symmetry is also behind the remarkable reduction in the number of distinct frequencies we noted at the end of Sec. 4, from a possible $N(N+1)/2$ to only five distinct frequencies.

Parenthetically, we should remark here that $\mathbf{G}$ and $\mathbf{F}$ are also symmetric matrices ($\mathbf{G}^T = \mathbf{G}$ and $\mathbf{F}^T = \mathbf{F}$); however, while $\mathbf{FG}$ does display the high degree of symmetry of Eqs. (57), it is not a symmetric matrix.

5.3 $\mathbf{Q}$ matrices in terms of simple submatrices

The S_N symmetry of the $\mathbf{Q}$ matrices ($\mathbf{F}$, $\mathbf{G}$, and $\mathbf{FG}$) described in Eqs. (57) allows us to write these matrices in terms of six simple submatrices. We first define the number of $\overline{\gamma}_{ij}$ coordinates to be

$$M \equiv N(N-1)/2, \quad (58)$$

and let $\mathbf{I}_N$ be an $N \times N$ identity matrix, $\mathbf{I}_M$ an $M \times M$ identity matrix, $\mathbf{J}_N$ an $N \times N$ matrix of ones and $\mathbf{J}_M$ an $M \times M$ matrix of ones. Further, we let $\mathbf{R}$ be an $N \times M$ matrix such that $R_{i,jk} = \delta_{ij} + \delta_{ik}$, $\mathbf{J}_{NM}$ be an $N \times M$ matrix of ones, and $\mathbf{J}_{NM}^T = \mathbf{J}_{MN}$.

We then write the $\mathbf{Q}$ matrices as

$$\mathbf{Q} = \begin{pmatrix} \mathbf{Q}_{\overline{r}'\overline{r}'} & \mathbf{Q}_{\overline{r}'\overline{\gamma}'} \\ \mathbf{Q}_{\overline{\gamma}'\overline{r}'} & \mathbf{Q}_{\overline{\gamma}'\overline{\gamma}'} \end{pmatrix}, \quad (59)$$

where the block $\mathbf{Q}_{\overline{r}'\overline{r}'}$ has dimension $(N \times N)$, block $\mathbf{Q}_{\overline{r}'\overline{\gamma}'}$ has dimension $(N \times M)$, block $\mathbf{Q}_{\overline{\gamma}'\overline{r}'}$ has dimension $(M \times N)$, and block $\mathbf{Q}_{\overline{\gamma}'\overline{\gamma}'}$ has dimension $(M \times M)$. Now, as we show in Appendix B of Ref. [2], Eqs. (57) allows us to write the following:

$$\mathbf{Q}_{\overline{r}'\overline{r}'} = (Q_a - Q_b)\mathbf{I}_N + Q_b\mathbf{J}_N \quad (60)$$

$$\mathbf{Q}_{\overline{r}'\overline{\gamma}'} = (Q_e - Q_f)\mathbf{R} + Q_f\mathbf{J}_{NM} \quad (61)$$

$$\mathbf{Q}_{\overline{\gamma}'\overline{r}'} = (Q_c - Q_d)\mathbf{R}^T + Q_d\mathbf{J}_{NM}^T \quad (62)$$

$$\mathbf{Q}_{\overline{\gamma}'\overline{\gamma}'} = (Q_g - 2Q_h + Q_\iota)\mathbf{I}_M + (Q_h - Q_\iota)\mathbf{R}^T\mathbf{R} + Q_\iota\mathbf{J}_M. \quad (63)$$

In particular, letting $\mathbf{Q} = \mathbf{F}\mathbf{G}$, the matrix that must be diagonalized, Eq. (59) becomes

$$\mathbf{F}\mathbf{G} = \begin{pmatrix} \tilde{a}\mathbf{I}_N + \tilde{b}\mathbf{J}_N & \tilde{e}\mathbf{R} + \tilde{f}\mathbf{J}_{NM} \\ \tilde{c}\mathbf{R}^T + \tilde{d}\mathbf{J}_{MN} & \tilde{g}\mathbf{I}_M + \tilde{h}\mathbf{R}^T\mathbf{R} + \tilde{l}\mathbf{J}_M \end{pmatrix}, \quad (64)$$

where we have used the following abbreviations:

$$\begin{aligned} \tilde{a} &\equiv (FG)_a - (FG)_b = (F_a - F_b)G_a \\ \tilde{b} &\equiv (FG)_b = F_b G_a \\ \tilde{c} &\equiv (FG)_c - (FG)_d = (F_e - F_f)G_a \\ \tilde{d} &\equiv (FG)_d = F_f G_a \\ \tilde{e} &\equiv (FG)_e - (FG)_f = (F_e - F_f)(G_g + (N-4)G_h) \\ \tilde{f} &\equiv (FG)_f = 2F_e G_h + F_f(G_g + 2(N-3)G_h) \\ \tilde{g} &\equiv (FG)_g - 2(FG)_h + (FG)_\iota = (F_g - 2F_h + F_\iota)(G_g - 2G_h) \\ \tilde{h} &\equiv (FG)_h - (FG)_\iota = F_g G_h + F_h(G_g + (N-6)G_h) - F_\iota(G_g + (N-5)G_h) \\ \tilde{l} &\equiv (FG)_\iota = 4F_h G_h + F_\iota(G_g + 2(N-4)G_h). \end{aligned} \quad (65)$$

The right-hand sides of Eq. (65), the $\mathbf{F}\mathbf{G}$ matrices expressed in terms of the $\mathbf{F}$ and $\mathbf{G}$ matrix elements, may be derived using the graph-theoretic techniques discussed in Appendix B of Ref. [2].

We also require the $\mathbf{G}$ matrix for the normalization condition (Eq. (49)). It has a simpler structure than the $\mathbf{F}\mathbf{G}$ matrix,

$$\mathbf{G} = \begin{pmatrix} \tilde{a}'\mathbf{I}_N & \mathbf{0} \\ \mathbf{0} & \tilde{g}'\mathbf{I}_M + \tilde{h}'\mathbf{R}^T\mathbf{R} \end{pmatrix}, \quad (66)$$

where

$$\begin{aligned} \tilde{a}' &\equiv (G)_a \\ \tilde{g}' &\equiv (G)_g - 2(G)_h \\ \tilde{h}' &\equiv (G)_h \end{aligned} \quad (67)$$

and

$$\begin{aligned} (G)_b &= 0 \\ (G)_c &= 0 \\ (G)_d &= 0 \\ (G)_e &= 0 \\ (G)_f &= 0 \\ (G)_\iota &= 0. \end{aligned} \quad (68)$$

The quantities $(G)_a$, $(G)_g$ and $(G)_h$ depend on the choice of $\kappa(D)$ (See Ref. [2]).

6 Symmetry and Normal Coordinates

The $\mathbf{FG}$ matrix is a $N(N+1)/2 \times N(N+1)/2$ dimensional matrix (there being $N(N+1)/2$ internal coordinates), and so Eqs. (47) and (48) could have up to $N(N+1)/2$ distinct frequencies. However, as noted above, there are only five distinct frequencies. The S_N symmetry is responsible for the remarkable reduction from $N(N+1)/2$ possible distinct frequencies to five actual distinct frequencies. As we shall also see, the S_N symmetry greatly simplifies the determination of the normal coordinates and hence the solution of the large- D problem.

6.1 Symmetrized coordinates

To understand why there are only five distinct frequencies and to start the process of solving for the normal coordinates, we first look at the S_N transformation properties of the internal coordinates.

We first note that the internal-coordinate displacement vectors $\bar{\mathbf{r}}'$ and $\bar{\boldsymbol{\gamma}}'$ of Eqs. (39) and (40) are basis functions which transform under matrix representations of S_N and each span the corresponding carrier spaces. These representations of S_N however, are not irreducible representations of S_N (Appendix C).

We will show that the reducible representation under which $\bar{\mathbf{r}}'$ transforms is reducible to one 1-dimensional irreducible representation labelled by the partition $[N]$ (the partition denotes a corresponding Young diagram (= Young pattern = Young shape) of an irreducible representation (see Appendix C)) and one $(N-1)$ -dimensional irreducible representation labelled by the partition $[N-1, 1]$. We will also show that the reducible representation under which $\bar{\boldsymbol{\gamma}}'$ transforms is reducible to one 1-dimensional irreducible representation labelled by the partition $[N]$, one $(N-1)$ -dimensional irreducible representation labelled by the partition $[N-1, 1]$ and one $N(N-3)/2$ -dimensional irreducible representation labelled by the partition $[N-2, 2]$.

6.1.1 The Characters of the Reducible Representations of S_N under which $\bar{\mathbf{r}}'$ and $\bar{\boldsymbol{\gamma}}'$ Transform.

Let us denote the cycle structure of a permutation (see Appendix C) by the symbol $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$, where the notation j^{ν_j} means a cycle of length j and ν_j equals the number of cycles of length j in that permutation[15]. Crucially the characters, $\chi^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}$, for all elements of a matrix representation of S_N corresponding to permutations with the same cycle structure $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$, are identical (Appendix C).

We will use the theory of group characters to decompose $\bar{\mathbf{r}}'$ and $\bar{\boldsymbol{\gamma}}'$ into basis functions which transform under irreducible representations of S_N . The theory of group characters is briefly discussed in Appendix C. To determine the characters, $\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}$, of the representation under which $\bar{\mathbf{r}}'$ transforms we need to calculate how many of the elements $\bar{\mathbf{r}}'$ transform into themselves under a permutation in the class $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$. As any number occurring in a cycle of length greater than one is transformed into another number, only those elements of $\bar{\mathbf{r}}'$, r_i , for which the index i occurs in a cycle of length one transform

into themselves. As there are ν_1 cycles of length one, there will also be ν_1 elements of $\bar{\mathbf{r}}'$ which transform into themselves. Hence the character of the representation under which $\bar{\mathbf{r}}'$ transforms under a permutation in the class $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$ is given by

$$\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} = \nu_1. \quad (69)$$

The derivation of the characters, $\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}$, under which the $\bar{\mathbf{r}}'$ transforms is similar. For a given permutation in the class $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$ all pairs of indices occurring in cycle of length one correspond to an element of $\bar{\mathbf{r}}'$, $\bar{r}'_{ij}$, for which the indices i or j transform into themselves. There will be $\nu_1(\nu_1 - 1)/2$ such pairs. Also every pair of indices occurring in a cycle of length two correspond to an element of $\bar{\mathbf{r}}'$, $\bar{r}'_{ij}$, for which the indices i and j transform into themselves. Thus the element transforms into itself. There are ν_2 such pairs. Thus the $\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}$, under which the $\bar{\mathbf{r}}'$ transforms is the sum of these two numbers, i.e.

$$\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} = \frac{\nu_1(\nu_1 - 1)}{2} + \nu_2. \quad (70)$$

6.1.2 The Characters of the Irreducible Representations of S_N .

Now let's turn to the characters of the irreducible representations. It can be shown[15] that for an element of an irreducible matrix representation which corresponds to a particular permutation with cycle structure $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$, the characters of the $[N]$, $[N - 1, 1]$ and $[N - 2, 2]$ representations are:

$$\begin{aligned} \chi_{[N]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} &= 1 \\ \chi_{[N-1, 1]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} &= \nu_1 - 1 \\ \chi_{[N-2, 2]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} &= \frac{(\nu_1 - 1)(\nu_1 - 2)}{2} + \nu_2 - 1. \end{aligned} \quad (71)$$

Armed with these characters, we can now see which irreducible representations of S_N $\bar{\mathbf{r}}'$ and $\bar{\mathbf{r}}'$ span. In Appendix C we note that the characters of the representations under which $\bar{\mathbf{r}}'$ and $\bar{\mathbf{r}}'$ transform can be written as

$$\chi(R) = \sum_{\alpha} a_{\alpha} \chi_{\alpha}(R), \quad (72)$$

where R denotes the element of the group, $\chi(R)$ is the character of the representation under which $\bar{\mathbf{r}}'$ or $\bar{\mathbf{r}}'$ transform, α labels the irreducible representation of S_N , and $\chi_{\alpha}(R)$ is the character of the irreducible representation denoted by α . The decomposition of Eq. (72) is unique where the coefficient a_{α} is the number of times the irreducible representation labelled by α appears in the decomposition of the representation under which $\bar{\mathbf{r}}'$ or $\bar{\mathbf{r}}'$ transform.

Thus the $\bar{\mathbf{r}}'$ vector must span a $[N]$ and a $[N - 1, 1]$ irreducible representation as from Eqs. (69) and (71)

$$\chi_{\bar{\mathbf{r}}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} = 1 \times \chi_{[N]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} + 1 \times \chi_{[N-1, 1]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}, \quad (73)$$

i.e.

$$a_{[N]} = 1 \quad (74)$$

$$a_{[N-1, 1]} = 1. \quad (75)$$

The $\overline{\gamma}'$ vector spans a $[N]$, a $[N-1, 1]$ and a $[N-2, 2]$ irreducible representation as from Eqs. (70) and (71)

$$\chi_{\overline{\gamma}'}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} = 1 \times \chi_{[N]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} + 1 \times \chi_{[N-1, 1]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})} + 1 \times \chi_{[N-2, 2]}^{(1^{\nu_1}, 2^{\nu_2}, \dots, N^{\nu_N})}, \quad (76)$$

i.e.

$$a_{[N]} = 1 \quad (77)$$

$$a_{[N-1, 1]} = 1 \quad (78)$$

$$a_{[N-2, 2]} = 1. \quad (79)$$

Regarding the dimensionalities of the $[N]$, $[N-1, 1]$ and $[N-2, 2]$ irreducible representations there are many rules to determine these. The best known is that the dimensionality of an irreducible representation of S_N is equal to the number of standard Young tableaux that the Young diagram for the irreducible representation allows. Perhaps the simplest rule however[17], is to fill the boxes of the Young diagram with numbers determined as follows. To the number of boxes in the row to the right of a box add the number of boxes in the column below the same box and then add one for the box itself. Once all of the boxes of the Young tableau have been filled with a number in this fashion, multiply all of the numbers in the boxes together. If we denote this product by N_T , then the dimensionality, d , of the representation is

$$d = \frac{N!}{N_T} \quad (80)$$

from which we deduce that

$$\begin{aligned} d_{[N]} &= \frac{N!}{N!} = 1 \\ d_{[N-1, 1]} &= \frac{N!}{[N!/(N-1)]} = N-1 \\ d_{[N-2, 2]} &= \frac{N!}{[2N!/(N(N-3))]} = \frac{N(N-3)}{2}. \end{aligned} \quad (81)$$

We note that

$$d_{[N]} + d_{[N-1, 1]} = N, \quad (82)$$

giving correctly the dimension of the $\overline{\mathbf{r}}'$ vector and that

$$d_{[N]} + d_{[N-1, 1]} + d_{[N-2, 2]} = \frac{N(N-1)}{2}, \quad (83)$$

giving correctly the dimension of the $\overline{\gamma}'$ vector.

6.1.3 The Reduction of $D_{\bar{\mathbf{r}}'}(\mathbf{R})$ by $W_{\bar{\mathbf{r}}'}$.

Let $D_{\bar{\mathbf{r}}'}(R)$ be the matrix representation of S_N under which the $\bar{\mathbf{r}}'$ vector transforms. The considerations of Secs. 6.1.1 and 6.1.2 then imply that $D_{\bar{\mathbf{r}}'}(R)$ may be block diagonalized under a similarity transformation to

$$D_{S_{\bar{\mathbf{r}}'}}(R) = \left(\begin{array}{c|c} D_{S_{\bar{\mathbf{r}}'}}^{[N]}(R) & \mathbf{0} \\ \hline \mathbf{0} & D_{S_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R) \end{array} \right), \quad (84)$$

where $D_{S_{\bar{\mathbf{r}}'}}^{[N]}(R)$ is the one-dimensional irreducible matrix representation $[N]$ of S_N and $D_{S_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R)$ is an $(N-1) \times (N-1)$ -dimensional irreducible matrix representation $[N-1, 1]$ of S_N .

If $W_{\bar{\mathbf{r}}'}$ is a matrix which effects the reduction of $D_{\bar{\mathbf{r}}'}(R)$ to Eq. (84), then $W_{\bar{\mathbf{r}}'}$ satisfies the equation

$$W_{\bar{\mathbf{r}}'} D_{\bar{\mathbf{r}}'}(R) W_{\bar{\mathbf{r}}'}^{-1} [W_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}'] = D_{S_{\bar{\mathbf{r}}'}}(R) [W_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}']. \quad (85)$$

Thus we have that the symmetry coordinates, $\mathbf{S}_{\bar{\mathbf{r}}'}$, of the $\bar{\mathbf{r}}'$ sector which transform under irreducible representations of the group S_N are given by

$$\mathbf{S}_{\bar{\mathbf{r}}'} = W_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}' \quad (86)$$

and so writing

$$W_{\bar{\mathbf{r}}'} = \begin{pmatrix} W_{\bar{\mathbf{r}}'}^{[N]} \\ W_{\bar{\mathbf{r}}'}^{[N-1, 1]} \end{pmatrix}, \quad (87)$$

where $W_{\bar{\mathbf{r}}'}^{[N]}$ is a $1 \times N$ dimensional matrix and $W_{\bar{\mathbf{r}}'}^{[N-1, 1]}$ is an $(N-1) \times N$ dimensional matrix, then

$$\mathbf{S}_{\bar{\mathbf{r}}'} = \begin{pmatrix} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]} \\ \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]} \end{pmatrix} = \begin{pmatrix} W_{\bar{\mathbf{r}}'}^{[N]} \bar{\mathbf{r}}' \\ W_{\bar{\mathbf{r}}'}^{[N-1, 1]} \bar{\mathbf{r}}' \end{pmatrix}. \quad (88)$$

The symmetry coordinate column vector $\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}$ is a one-element vector and transforms under $D_{S_{\bar{\mathbf{r}}'}}^{[N]}(R)$, the one-dimensional irreducible matrix representation $[N]$ of S_N . The symmetry coordinate column vector $\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]}$ is a $(N-1)$ -element vector and transforms under $D_{S_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R)$, the $(N-1) \times (N-1)$ -dimensional irreducible matrix representation $[N-1, 1]$ of S_N .

6.1.4 Reduction of $D_{\bar{\mathbf{r}}'}(\mathbf{R})$ by $W_{\bar{\mathbf{r}}'}$

. Likewise, if $W_{\bar{\mathbf{r}}'}$ is a matrix which effects the reduction of the representation, $D_{\bar{\mathbf{r}}'}(R)$, under which the $\bar{\mathbf{r}}'$ vector transforms then

$$W_{\bar{\mathbf{r}}'} D_{\bar{\mathbf{r}}'}(R) W_{\bar{\mathbf{r}}'}^{-1} [W_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}'] = D_{S_{\bar{\mathbf{r}}'}}(R) [W_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}'], \quad (89)$$

where

$$D_{S_{\bar{\mathbf{r}}'}}(R) = \left(\begin{array}{c|c|c} D_{S_{\bar{\mathbf{r}}'}}^{[N]}(R) & \mathbf{0} & \mathbf{0} \\ \hline \mathbf{0} & D_{S_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R) & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & D_{S_{\bar{\mathbf{r}}'}}^{[N-2, 2]}(R) \end{array} \right), \quad (90)$$

$D_{S_{\vec{\gamma}'}}^{[N]}(R)$ is the one-dimensional irreducible matrix representation $[N]$ of S_N , $D_{S_{\vec{\gamma}'}}^{[N-1, 1]}(R)$ is the $(N-1) \times (N-1)$ -dimensional irreducible matrix representation $[N-1, 1]$ of S_N , and $D_{S_{\vec{\gamma}'}}^{[N-2, 2]}(R)$ is the $\{N(N-3)/2\} \times \{N(N-3)/2\}$ -dimensional irreducible matrix representation $[N-2, 2]$ of S_N .

Note that while $D_{\vec{\gamma}'}^\alpha(R)$ and $D_{\vec{\gamma}}^\alpha(R)$ both belong to the α irreducible representation of S_N , where α is the partition $[N]$ or $[N-1, 1]$, they are not necessarily the same matrices. Generally they are related by an equivalence transformation

$$D_{\vec{\gamma}'}^\alpha(R) = t D_{\vec{\gamma}}^\alpha(R) t^{-1}, \quad (91)$$

where t is the transformation matrix.

Thus we have that the symmetry coordinates, $\mathbf{S}_{\vec{\gamma}'}$, of the $\vec{\gamma}'$ sector which transform under irreducible representations of the group S_N are given by

$$\mathbf{S}_{\vec{\gamma}'} = W_{\vec{\gamma}'} \vec{\gamma}' \quad (92)$$

and so writing

$$W_{\vec{\gamma}'} = \begin{pmatrix} W_{\vec{\gamma}'}^{[N]} \\ W_{\vec{\gamma}'}^{[N-1, 1]} \\ W_{\vec{\gamma}'}^{[N-2, 2]} \end{pmatrix}, \quad (93)$$

where $W_{\vec{\gamma}'}^{[N]}$ is a $1 \times N(N-1)/2$ dimensional matrix, $W_{\vec{\gamma}'}^{[N-1, 1]}$ is an $(N-1) \times N(N-1)/2$ dimensional matrix and $W_{\vec{\gamma}'}^{[N-2, 2]}$ is an $N(N-3)/2 \times N(N-1)/2$ dimensional matrix, then

$$\mathbf{S}_{\vec{\gamma}'} = \begin{pmatrix} \mathbf{S}_{\vec{\gamma}'}^{[N]} \\ \mathbf{S}_{\vec{\gamma}'}^{[N-1, 1]} \\ \mathbf{S}_{\vec{\gamma}'}^{[N-2, 2]} \end{pmatrix} = \begin{pmatrix} W_{\vec{\gamma}'}^{[N]} \vec{\gamma}' \\ W_{\vec{\gamma}'}^{[N-1, 1]} \vec{\gamma}' \\ W_{\vec{\gamma}'}^{[N-2, 2]} \vec{\gamma}' \end{pmatrix}, \quad (94)$$

where

$$W_{\vec{\gamma}'}^\alpha \vec{\gamma}' = \sum_{j=1}^N \sum_{i < j} [W_{\vec{\gamma}'}^\alpha]_{ij} \vec{\gamma}'_{ij}. \quad (95)$$

The symmetry coordinate column vector $\mathbf{S}_{\vec{\gamma}'}^{[N]}$ is a one-element vector and transforms under $D_{S_{\vec{\gamma}'}}^{[N]}(R)$, the one-dimensional irreducible matrix representation $[N]$ of S_N , while the symmetry coordinate column vector $\mathbf{S}_{\vec{\gamma}'}^{[N-1, 1]}$ is an $(N-1)$ -element vector and transforms under $D_{S_{\vec{\gamma}'}}^{[N-1, 1]}(R)$, the $(N-1) \times (N-1)$ -dimensional irreducible matrix representation $[N-1, 1]$ of S_N . The symmetry coordinate column vector $\mathbf{S}_{\vec{\gamma}'}^{[N-2, 2]}$ is an $N(N-3)/2$ -element vector and transforms under $D_{S_{\vec{\gamma}'}}^{[N-2, 2]}(R)$, the $\{N(N-3)/2\} \times \{N(N-3)/2\}$ -dimensional irreducible matrix representation $[N-2, 2]$ of S_N .

6.1.5 The Full Symmetry Coordinate Vector $\mathbf{S}$.

Symmetry coordinates $\mathbf{S}_{\tilde{r}'}^\alpha$ and $\mathbf{S}_{\tilde{\gamma}'}^\beta$ are said to belong to the same species when $\alpha = \beta$. Now in what follows we will find it useful to form a full symmetry coordinate vector as follows:

$$\mathbf{S} = P \begin{pmatrix} \mathbf{S}_{\tilde{r}'} \\ \mathbf{S}_{\tilde{\gamma}'} \end{pmatrix} = \begin{pmatrix} \mathbf{S}_{\tilde{r}'}^{[N]} \\ \mathbf{S}_{\tilde{\gamma}'}^{[N]} \\ \mathbf{S}_{\tilde{r}'}^{[N-1, 1]} \\ \mathbf{S}_{\tilde{\gamma}'}^{[N-1, 1]} \\ \mathbf{S}_{\tilde{r}'}^{[N-2, 2]} \\ \mathbf{S}_{\tilde{\gamma}'}^{[N-2, 2]} \end{pmatrix} = \begin{pmatrix} \mathbf{S}^{[N]} \\ \mathbf{S}^{[N-1, 1]} \\ \mathbf{S}^{[N-2, 2]} \end{pmatrix}, \quad (96)$$

where the orthogonal matrix

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (97)$$

and

$$\mathbf{S}^{[N]} = \begin{pmatrix} \mathbf{S}_{\tilde{r}'}^{[N]} \\ \mathbf{S}_{\tilde{\gamma}'}^{[N]} \end{pmatrix}, \quad \mathbf{S}^{[N-1, 1]} = \begin{pmatrix} \mathbf{S}_{\tilde{r}'}^{[N-1, 1]} \\ \mathbf{S}_{\tilde{\gamma}'}^{[N-1, 1]} \end{pmatrix} \quad \text{and} \quad \mathbf{S}^{[N-2, 2]} = \mathbf{S}_{\tilde{\gamma}'}^{[N-2, 2]}. \quad (98)$$

In $\mathbf{S}$, symmetry coordinates of the same species are grouped together.

6.2 Normal Coordinates

So how do symmetry coordinates simplify the solution of Eqs. (46), (47), (48), (49) and (52) for the frequencies and normal modes? Symmetry coordinates have maximum utility when

1. all transformation matrices t , which relate the equivalent irreducible representations of the symmetry group under which symmetry coordinates of the same species transform (see Eq. (91)), are equal to the identity matrix. The irreducible representations are not just equivalent, they are identical.
2. the symmetry coordinates transform under unitary representations of the symmetry group (here S_N). In our case we wish to have real symmetry coordinates and normal mode coordinates, and so we require that the symmetry coordinates transform under orthogonal representations of S_N .

Now consider applying a transformation W to Eqs. (46), (47), (48) and (49), where

$$W = P \left(\begin{array}{c|c} W_{\tilde{r}'} & \mathbf{0} \\ \hline \mathbf{0} & W_{\tilde{\gamma}'} \end{array} \right) = \begin{pmatrix} W_{\tilde{r}'}^{[N]} & \mathbf{0} \\ \mathbf{0} & W_{\tilde{\gamma}'}^{[N]} \\ \hline W_{\tilde{r}'}^{[N-1, 1]} & \mathbf{0} \\ \mathbf{0} & W_{\tilde{\gamma}'}^{[N-1, 1]} \\ \hline \mathbf{0} & W_{\tilde{\gamma}'}^{[N-2, 2]} \end{pmatrix}, \quad (99)$$

where

$$\mathbf{S} = W \bar{\mathbf{y}}'. \quad (100)$$

6.2.1 The Essential Equation and the Orthogonality of W .

In Appendix E we show that condition 2 above implies that W has to be an orthogonal matrix, i.e. W satisfies

$$WW^T = W^TW = I. \quad (101)$$

In particular we show that W is an orthogonal matrix when

$$W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^\alpha]^T = I_\alpha, \quad (102)$$

where $\mathbf{X}'$ is $\bar{\mathbf{r}}'$ or $\bar{\boldsymbol{\gamma}}'$ and I_α is the unit matrix.

Thus Eq. (102) is the *essential equation* for $W_{\mathbf{X}'}^\alpha$ to satisfy.

From Eq. (99)

$$W^TW = \begin{pmatrix} \sum_{\alpha=[N]}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^\alpha]^T W_{\bar{\mathbf{r}}'}^\alpha & \mathbf{0} \\ \mathbf{0} & \sum_{\alpha=[N]}^{[N-2, 2]} [W_{\bar{\boldsymbol{\gamma}}'}^\alpha]^T W_{\bar{\boldsymbol{\gamma}}'}^\alpha \end{pmatrix} \quad (103)$$

which with Eq. (101) gives us

$$\sum_{\alpha} [W_{\mathbf{X}'}^\alpha]^T W_{\mathbf{X}'}^\alpha = I_{\mathbf{X}'}, \quad (104)$$

where $I_{\mathbf{X}'}$ is the $N \times N$ -dimensional unit matrix when $\mathbf{X}' = \bar{\mathbf{r}}'$ or the $N(N-1)/2 \times N(N-1)/2$ -dimensional unit matrix when $\mathbf{X}' = \bar{\boldsymbol{\gamma}}'$. From Eq. (102) and the automatically satisfied Eq. (D4) we find

$$[W_{\mathbf{X}'}^{\alpha'}]^T W_{\mathbf{X}'}^{\alpha'} [W_{\mathbf{X}'}^\alpha]^T W_{\mathbf{X}'}^\alpha = \delta_{\alpha, \alpha'} [W_{\mathbf{X}'}^\alpha]^T W_{\mathbf{X}'}^\alpha, \quad (105)$$

and so Eq. (104) is a decomposition of the identity into primitive idempotent projection operators which annul each other on both sides. The $[W_{\mathbf{X}'}^\alpha]^T W_{\mathbf{X}'}^\alpha$ project out orthogonal irreducible subspaces while the $W_{\mathbf{X}'}^\alpha$ are the Clebsch-Gordon coefficients of this decomposition.

6.2.2 The Motion Associated with the Symmetry Coordinates about the Lewis Structure Configuration.

Equation (104) can be written as

$$I_{\mathbf{X}'} = \sum_{\alpha} \sum_{\xi} [(W_{\mathbf{X}'}^\alpha)_\xi]^T (W_{\mathbf{X}'}^\alpha)_\xi, \quad \text{where } \xi = 1 \text{ when } \alpha = [N], \quad (106)$$

$$1 \leq \xi \leq N-1 \text{ when } \alpha = [N-1, 1],$$

$$1 \leq \xi \leq N(N-3)/2 \text{ when } \alpha = [N-2, 2],$$

and the symbol ξ denotes the ξ^{th} row of $W_{\mathbf{X}'}$. Thus we can write

$$\mathbf{X}' = I_{\mathbf{X}'} \mathbf{X}' = \sum_{\alpha} [W_{\mathbf{X}'}^{\alpha}]^T (W_{\mathbf{X}'}^{\alpha} \cdot \mathbf{X}') = \sum_{\alpha} \sum_{\xi} ((W_{\mathbf{X}'}^{\alpha})_{\xi} \cdot \mathbf{X}') [(W_{\mathbf{X}'}^{\alpha})_{\xi}]^T, \quad (107)$$

where $\mathbf{X}'$ is $\bar{\mathbf{r}}'$ or $\bar{\boldsymbol{\gamma}}'$ of Eqs. (39) or (40) respectively. According to Eqs. (D4) and (102) the $[(W_{\mathbf{X}'}^{\alpha})_{\xi}]^T$ are column vectors which form a complete set of orthonormal basis functions where, from Eqs. (D4) (102), one has

$$(W_{\mathbf{X}'}^{\alpha})_{\xi} [(W_{\mathbf{X}'}^{\alpha'})^T]_{\xi'} = \delta_{\alpha, \alpha'} \delta_{\xi, \xi'}. \quad (108)$$

From Eqs. (88) and (94)

$$[\mathbf{S}_{\mathbf{X}'}^{\alpha}]_{\xi} = (W_{\mathbf{X}'}^{\alpha})_{\xi} \cdot \mathbf{X}' \quad (109)$$

and so Eq. (107) is a decomposition of the $\bar{\mathbf{r}}'$ and $\bar{\boldsymbol{\gamma}}'$ vectors in terms of the symmetry coordinates, i.e.

$$\mathbf{X}' = \sum_{\alpha} \sum_{\xi} \mathbf{X}'_{\xi}{}^{\alpha}, \quad (110)$$

where

$$\mathbf{X}'_{\xi}{}^{\alpha} = [\mathbf{S}_{\mathbf{X}'}^{\alpha}]_{\xi} [(W_{\mathbf{X}'}^{\alpha})_{\xi}]^T \quad (111)$$

expresses the motion associated with the symmetry coordinate $[\mathbf{S}_{\mathbf{X}'}^{\alpha}]_{\xi}$ in the original internal displacement coordinate, $\mathbf{X}'$ ($\bar{\mathbf{r}}'$ or $\bar{\boldsymbol{\gamma}}'$). Equation (110) can be written concisely as

$$\bar{\mathbf{y}}' = W^T \mathbf{S}, \quad (112)$$

where $\bar{\mathbf{y}}'$ is defined in Eqs. (38), (39) and (40), $\mathbf{S}$ is given by Eqs. (96), (97) and (98) while W is given by Eq. (99). Equation (112) may be directly derived from Eq. (100) using the orthogonality of the W matrix (Eq. (101)).

Using Eqs. (110) and (111) in Eqs. (22), (32) and (33) we find that

$$\mathbf{y} = \begin{pmatrix} \mathbf{r} \\ \boldsymbol{\gamma} \end{pmatrix} = \begin{pmatrix} D^2 \bar{a}_{ho} \left(\bar{\mathbf{r}}'_{\infty} \mathbf{1}_{\bar{\mathbf{r}}'} + \frac{1}{\sqrt{D}} \sum_{\alpha} \sum_{\xi} \bar{\mathbf{r}}'_{\xi}{}^{\alpha} \right) \\ \bar{\boldsymbol{\gamma}}'_{\infty} \mathbf{1}_{\bar{\boldsymbol{\gamma}}'} + \frac{1}{\sqrt{D}} \sum_{\alpha} \sum_{\xi} \bar{\boldsymbol{\gamma}}'_{\xi}{}^{\alpha} \end{pmatrix} \quad (113)$$

where according to Eq. (111)

$$\bar{\mathbf{r}}'_{\xi}{}^{\alpha} = [\mathbf{S}_{\bar{\mathbf{r}}'}^{\alpha}]_{\xi} [(W_{\bar{\mathbf{r}}'}^{\alpha})_{\xi}]^T, \quad (114)$$

$$\bar{\boldsymbol{\gamma}}'_{\xi}{}^{\alpha} = [\mathbf{S}_{\bar{\boldsymbol{\gamma}}'}^{\alpha}]_{\xi} [(W_{\bar{\boldsymbol{\gamma}}'}^{\alpha})_{\xi}]^T, \quad (115)$$

while

$$[\mathbf{1}_{\bar{\mathbf{r}}'}]_i = 1 \quad \forall \quad 1 \leq i \leq N \quad (116)$$

and

$$[\mathbf{1}_{\bar{\boldsymbol{\gamma}}'}]_{ij} = 1 \quad \forall \quad 1 \leq i, j \leq N. \quad (117)$$

Equations (113), (114), (115), (116) and (117) express the motion associated with the symmetry coordinate about the Lewis structure configuration.

6.3 The Reduction of the Eigensystem Equations in the Symmetry Coordinate Basis .

6.3.1 The Central Theorem

Let us start out by defining $\mathbf{Q}_W$ to be the $\mathbf{Q}$ matrix in the symmetry coordinate basis, i.e.

$$\mathbf{Q}_W = W\mathbf{Q}W^T, \quad (118)$$

where $\mathbf{Q}$ is $\mathbf{F}$, $\mathbf{G}$ or $\mathbf{FG}$, and $\mathbf{Q}_W$ is $\mathbf{F}_W$, $\mathbf{G}_W$ or $(\mathbf{FG})_W$. It has been shown elsewhere[18] that since $\mathbf{F}$, $\mathbf{G}$ and $\mathbf{FG}$ are invariant matrices (under S_N), when items 1 and 2 above are satisfied then

$$\mathbf{Q}_W = \begin{pmatrix} \sigma_{[N]}^Q \otimes \mathbf{I}_{[N]} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \sigma_{[N-1, 1]}^Q \otimes \mathbf{I}_{[N-1, 1]} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \sigma_{[N-2, 2]}^Q \otimes \mathbf{I}_{[N-2, 2]} \end{pmatrix}, \quad (119)$$

where the symbol $\otimes$ indicates the direct product, $\sigma_{[N]}^Q$ and $\sigma_{[N-1, 1]}^Q$ are 2×2 dimensional matrices and $\sigma_{[N-2, 2]}^Q$ is a single element (a number). The matrix $\mathbf{I}_{[N]}$ is the $[N]$ -sector identity matrix, simply the number 1, $\mathbf{I}_{[N-1, 1]}$ is the $[N-1, 1]$ -sector identity matrix, the $(N-1) \times (N-1)$ -dimensional unit matrix and $\mathbf{I}_{[N-2, 2]}$ is the $[N-2, 2]$ -sector identity matrix, the $N(N-3)/2 \times N(N-3)/2$ -dimensional unit matrix. The matrix elements

$$[\sigma_\alpha^Q]_{\mathbf{X}'_1, \mathbf{X}'_2} = (W_{\mathbf{X}'_1}^\alpha)_\xi \mathbf{Q}_{\mathbf{X}'_1 \mathbf{X}'_2} [(W_{\mathbf{X}'_2}^\alpha)_\xi]^T, \quad (120)$$

where $\mathbf{X}'_1$ and $\mathbf{X}'_2$ are $\bar{r}'$ or $\bar{\gamma}'$ (only $\bar{\gamma}'$ for the $[N-2, 2]$ sector). The $\mathbf{Q}$ -matrix quadrant, $\mathbf{Q}_{\mathbf{X}'_1 \mathbf{X}'_2}$, is given by Eqs. (59), (60), (61), (62) and (63). Note that although we are *not* summing over the repeated index ξ in Eq. (120), the $[\sigma_\alpha^Q]_{\mathbf{X}'_1, \mathbf{X}'_2}$ are *independent* of the $W_{\mathbf{X}'}^\alpha$, row label, ξ . If, when we calculate $W_{\mathbf{X}'}^\alpha$, the matrix element $[\sigma_\alpha^Q]_{\mathbf{X}'_1, \mathbf{X}'_2}$ turns out to depend on the $W_{\mathbf{X}'}^\alpha$, row label, ξ , then we know that we have made a mistake calculating $W_{\mathbf{X}'}^\alpha$. This is a strong check on the correctness of our calculations in Sect. 7.

6.3.2 The Reduction of the Eigenvalue Equation in the Symmetry Coordinate Basis .

Transforming the basic eigenvalue equation, Eq. (47), to the symmetry coordinate basis we have

$$W\mathbf{F}W^T W\mathbf{G}W^T W\mathbf{b} = \mathbf{F}_W \mathbf{G}_W \mathbf{c}^{(b)} = \lambda_b \mathbf{c}^{(b)}, \quad (121)$$

where

$$\mathbf{F}_W = W\mathbf{F}W^T \quad \mathbf{G}_W = W\mathbf{G}W^T \quad \mathbf{c}^{(b)} = W\mathbf{b}. \quad (122)$$

Under this transformation we can also write Eq. (46) for the b^{th} normal-mode coordinate as

$$[q']_b = \mathbf{b}^T W^T W \bar{\mathbf{y}}' = [\mathbf{c}^{(b)}]^T \cdot \mathbf{S}, \quad (123)$$

where $[q']_b$ is now directly expressed in terms of the symmetry coordinates.

From Eqs. (119) and (121) the normal-coordinate coefficient vector, $\mathbf{c}^{(b)}$, has the form

$$\mathbf{c}^{(b)} = \begin{pmatrix} \delta_{\alpha, [N]} \mathbf{c}^{[N]} \otimes \mathbf{1} \\ \delta_{\alpha, [N-1, 1]} \mathbf{c}^{[N-1, 1]} \otimes \mathbf{1}_{\xi}^{[N-1, 1]} \\ \delta_{\alpha, [N-2, 2]} \mathbf{c}^{[N-2, 2]} \otimes \mathbf{1}_{\xi}^{[N-2, 2]} \end{pmatrix}, \quad (124)$$

where the $\mathbf{c}^{\alpha}$ satisfy the eigenvalue equations

$$\sigma_{\alpha}^{\mathbf{F}\mathbf{G}} \mathbf{c}^{\alpha} = \lambda_{\alpha} \mathbf{c}^{\alpha}. \quad (125)$$

Note that $\sigma_{[N]}^{\mathbf{F}\mathbf{G}}$ and $\sigma_{[N-1, 1]}^{\mathbf{F}\mathbf{G}}$ are 2×2 -dimensional matrices, while $\sigma_{[N-2, 2]}^{\mathbf{F}\mathbf{G}}$ is a one-dimensional matrix. Thus there are five solutions to Eq. (125) which we denote as $\{\lambda_{[N]}^{\pm}, \mathbf{c}_{\pm}^{[N]}\}$, $\{\lambda_{[N-1, 1]}^{\pm}, \mathbf{c}_{\pm}^{[N-1, 1]}\}$ and $\{\lambda_{[N-2, 2]}, \mathbf{c}^{[N-2, 2]}\}$, where $\mathbf{X}'$ ($\bar{\mathbf{r}}'$ or $\bar{\mathbf{r}}'$, only $\bar{\mathbf{r}}'$ for the $[N-2, 2]$ sector) labels the rows of the elements of the column vector $\mathbf{c}^{\alpha}$. The normal-coordinate label, b , has been replaced by the labels α , ξ and $\pm$ on the right hand side of Eq. (124), while the elements of the column vectors $\mathbf{1}_{\xi}^{\alpha}$ are

$$[\mathbf{1}_{\xi}^{\alpha}]_{\eta} = \delta_{\xi\eta}, \quad (126)$$

with $1 \leq \xi, \eta \leq N-1$ when $\alpha = [N-1, 1]$, or $1 \leq \xi, \eta \leq N(N-3)/2$ when $\alpha = [N-2, 2]$. The $\mathbf{c}_{\pm}^{\alpha}$ for the $[N]$ and $[N-1, 1]$ sectors determine the amount of angular-radial mixing between the symmetry coordinates in a normal coordinate of a particular α since from Eqs. (96), (98) and (123)

$$[q']_b = [\mathbf{c}_{\pm}^{\alpha}]_{\bar{\mathbf{r}}'} [\mathbf{S}_{\bar{\mathbf{r}}'}^{\alpha}]_{\xi} + [\mathbf{c}_{\pm}^{\alpha}]_{\bar{\mathbf{r}}'} [\mathbf{S}_{\bar{\mathbf{r}}'}^{\alpha}]_{\xi}. \quad (127)$$

For the $[N-2, 2]$ sector we have

$$[q']_b = [\mathbf{c}^{[N-2, 2]}]_{\bar{\mathbf{r}}'} [\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]}]_{\xi}, \quad (128)$$

i.e. the $[N-2, 2]$ sector symmetry coordinates are the $[N-2, 2]$ sector normal coordinates up to a normalization constant, $[\mathbf{c}^{[N-2, 2]}]_{\bar{\mathbf{r}}'}$ (see Sec. 6.3.3).

If we write $\mathbf{c}^{\alpha}$ as

$$\mathbf{c}^{\alpha} = c^{\alpha} \times \widehat{\mathbf{c}}^{\alpha} \quad (129)$$

where $\widehat{\mathbf{c}}^{\alpha}$ is a vector satisfying the normalization condition

$$[\widehat{\mathbf{c}}^{\alpha}]^T \widehat{\mathbf{c}}^{\alpha} = 1 \quad (130)$$

and c^{α} is a normalization factor ensuring that Eq. (49) is satisfied (see Sec. 6.3.3 below), then the reduced eigenvalue equation, Eq. (125), determines $\widehat{\mathbf{c}}^{\alpha}$ alone.

With the $[N-2, 2]$ sector, Eq. (130) yields

$$\widehat{\mathbf{c}^{[N-2, 2]}} = 1 \quad (131)$$

and so Eq. (125) yields directly

$$\lambda_{[N-2, 2]} = \sigma_{[N-2, 2]}^{\mathbf{F}\mathbf{G}}. \quad (132)$$

As for the $[N]$ and $[N - 1, 1]$ sectors, if we write

$$\widehat{\mathbf{c}}_{\pm}^{\alpha} = \begin{pmatrix} \cos \theta_{\pm}^{\alpha} \\ \sin \theta_{\pm}^{\alpha} \end{pmatrix}, \quad (133)$$

then from Eqs. (127), (129) and (133)

$$[q']_b = c_{\pm}^{\alpha} (\cos \theta_{\pm}^{\alpha} [\mathbf{S}_{\bar{r}'}^{\alpha}]_{\xi} + \sin \theta_{\pm}^{\alpha} [\mathbf{S}_{\bar{\gamma}'}^{\alpha}]_{\xi}). \quad (134)$$

Writing

$$\sigma_{\alpha}^{FG} = \begin{pmatrix} [\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{r}'} & [\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{\gamma}'} \\ [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{r}'} & [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} \end{pmatrix}, \quad (135)$$

then Eq. (125) may be written as

$$\begin{pmatrix} ([\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{r}'} - \lambda_{\alpha}^{\pm}) & [\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{\gamma}'} \\ [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{r}'} & ([\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} - \lambda_{\alpha}^{\pm}) \end{pmatrix} \begin{pmatrix} \cos \theta_{\pm}^{\alpha} \\ \sin \theta_{\pm}^{\alpha} \end{pmatrix} = 0, \quad (136)$$

from which we derive

$$\lambda_{\alpha}^{\pm} = \frac{([\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{r}'} + [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} \pm \sqrt{([\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{r}'} - [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{\gamma}'})^2 + 4[\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{\gamma}'} [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{r}'}}}{2} \quad (137)$$

Equations (136) and (137) then give us

$$\tan \theta_{\pm}^{\alpha} = \frac{(\lambda_{\alpha}^{\pm} - [\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{r}'})}{[\sigma_{\alpha}^{FG}]_{\bar{r}', \bar{\gamma}'}} = \frac{[\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{r}'}}{(\lambda_{\alpha}^{\pm} - [\sigma_{\alpha}^{FG}]_{\bar{\gamma}', \bar{\gamma}'})} \quad (138)$$

6.3.3 The Normalization Condition.

From Eqs. (49), (122), (119), (120) and (124) the $\mathbf{c}^{\alpha}$ also satisfy the normalization condition

$$[\mathbf{c}^{\alpha}]^T \sigma_{\alpha}^G \mathbf{c}^{\alpha} = 1. \quad (139)$$

Thus Eqs. (129) and (139) mean that

$$c^{\alpha} = \frac{1}{\sqrt{[\widehat{\mathbf{c}}^{\alpha}]^T \sigma_{\alpha}^G \widehat{\mathbf{c}}^{\alpha}}}. \quad (140)$$

For the $[N - 2, 2]$ sector, Eqs. (131) and (140) mean that

$$c^{[N-2, 2]} = \frac{1}{\sqrt{\sigma_{[N-2, 2]}^G}}. \quad (141)$$

As for the normalization constant, $c_{\pm}^{\alpha}$, of the $[N]$ and $[N - 1, 1]$ sectors, Eqs. (133) and (140) give us

$$c_{\pm}^{\alpha} = \frac{1}{\sqrt{\left(\begin{pmatrix} \cos \theta_{\pm}^{\alpha} \\ \sin \theta_{\pm}^{\alpha} \end{pmatrix}^T \sigma_{\alpha}^G \begin{pmatrix} \cos \theta_{\pm}^{\alpha} \\ \sin \theta_{\pm}^{\alpha} \end{pmatrix} \right)}}. \quad (142)$$

6.3.4 The Normal Coordinates.

Thus the total transformation matrix, V , from the internal displacement coordinates to the normal coordinates is

$$V = C W, \quad (143)$$

where W is given by Eq. (99) and

$$C = \left(\begin{array}{cc|cc|c} c_+^{[N]} \cos \theta_+^{[N]} & c_+^{[N]} \sin \theta_+^{[N]} & 0 & 0 & 0 \\ c_-^{[N]} \cos \theta_-^{[N]} & c_-^{[N]} \sin \theta_-^{[N]} & 0 & 0 & 0 \\ \hline 0 & 0 & c_+^{[N-1, 1]} \cos \theta_+^{[N-1, 1]} & c_+^{[N-1, 1]} \sin \theta_+^{[N-1, 1]} & 0 \\ 0 & 0 & c_-^{[N-1, 1]} \cos \theta_-^{[N-1, 1]} & c_-^{[N-1, 1]} \sin \theta_-^{[N-1, 1]} & 0 \\ \hline 0 & 0 & 0 & 0 & c^{[N-2, 2]} \end{array} \right), \quad (144)$$

and so the normal-coordinate vector, $\mathbf{q}'$, is given by

$$\begin{aligned} \mathbf{q}' &= \begin{pmatrix} \mathbf{q}'^{[N]} \\ \mathbf{q}'^{[N-1, 1]} \\ \mathbf{q}'^{[N-2, 2]} \end{pmatrix} = \begin{pmatrix} \frac{\mathbf{q}'_+^{[N]}}{\mathbf{q}'_-^{[N]}} \\ \frac{\mathbf{q}'_+^{[N-1, 1]}}{\mathbf{q}'_-^{[N-1, 1]}} \\ \frac{\mathbf{q}'_+^{[N-2, 2]}}{\mathbf{q}'_-^{[N-2, 2]}} \end{pmatrix} = V \bar{\mathbf{y}}' = C \mathbf{S} = C \begin{pmatrix} \frac{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}}{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}} \\ \frac{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]}}{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]}} \\ \frac{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]}}{\mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]}} \end{pmatrix} = \\ &= \begin{pmatrix} \frac{c_+^{[N]} \cos \theta_+^{[N]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]} + c_+^{[N]} \sin \theta_+^{[N]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}}{c_-^{[N]} \cos \theta_-^{[N]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]} + c_-^{[N]} \sin \theta_-^{[N]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}} \\ \frac{c_+^{[N-1, 1]} \cos \theta_+^{[N-1, 1]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]} + c_+^{[N-1, 1]} \sin \theta_+^{[N-1, 1]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]}}{c_-^{[N-1, 1]} \cos \theta_-^{[N-1, 1]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]} + c_-^{[N-1, 1]} \sin \theta_-^{[N-1, 1]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-1, 1]}} \\ \frac{c_+^{[N-2, 2]} \cos \theta_+^{[N-2, 2]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]} + c_+^{[N-2, 2]} \sin \theta_+^{[N-2, 2]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]}}{c_-^{[N-2, 2]} \cos \theta_-^{[N-2, 2]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]} + c_-^{[N-2, 2]} \sin \theta_-^{[N-2, 2]} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N-2, 2]}} \end{pmatrix}. \end{aligned} \quad (145)$$

6.3.5 The Motion Associated with the Normal Coordinates about the Lewis Structure Configuration.

Consider now the inverse transformation to express the internal displacement coordinates in terms of the normal coordinates. From Eq. (101)

$$I = W^T W = W^T C^{-1} C W \quad (146)$$

so that

$$\bar{\mathbf{y}}' = W^T C^{-1} (V \bar{\mathbf{y}}') = W^T C^{-1} \mathbf{q}', \quad (147)$$

where

$$C^{-1} = \left(\begin{array}{cc|cc|c} \frac{1}{s(\theta^{[N]})} \begin{pmatrix} -\frac{\sin \theta_-^{[N]}}{c_+^{[N]}} & \frac{\sin \theta_+^{[N]}}{c_-^{[N]}} \\ \frac{\cos \theta_-^{[N]}}{c_+^{[N]}} & -\frac{\cos \theta_+^{[N]}}{c_-^{[N]}} \end{pmatrix} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & \frac{1}{s(\theta^{[N-1, 1]})} \begin{pmatrix} -\frac{\sin \theta_-^{[N-1, 1]}}{c_+^{[N-1, 1]}} & \frac{\sin \theta_+^{[N-1, 1]}}{c_-^{[N-1, 1]}} \\ \frac{\cos \theta_-^{[N-1, 1]}}{c_+^{[N-1, 1]}} & -\frac{\cos \theta_+^{[N-1, 1]}}{c_-^{[N-1, 1]}} \end{pmatrix} & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & \frac{1}{c^{[N-2, 2]}} \end{array} \right) \quad (148)$$

and

$$s(\theta^\alpha) = \sin(\theta_+^\alpha - \theta_-^\alpha). \quad (149)$$

Thus from Eqs. (38), (147), (148) and (149)

$$\begin{aligned} \bar{\mathbf{y}}' = \begin{pmatrix} \bar{\mathbf{r}}' \\ \bar{\boldsymbol{\gamma}}' \end{pmatrix} = & \frac{1}{s(\theta^{[N]})} \begin{pmatrix} [W_{\bar{\mathbf{r}}'}^{[N]}]^T \left(\frac{-\sin \theta_-^{[N]}}{c_+^{[N]}} \mathbf{q}'_{+^{[N]}} + \frac{\sin \theta_+^{[N]}}{c_-^{[N]}} \mathbf{q}'_{-^{[N]}} \right) \\ [W_{\bar{\boldsymbol{\gamma}}'}^{[N]}]^T \left(\frac{\cos \theta_-^{[N]}}{c_+^{[N]}} \mathbf{q}'_{+^{[N]}} + \frac{-\cos \theta_+^{[N]}}{c_-^{[N]}} \mathbf{q}'_{-^{[N]}} \right) \end{pmatrix} + \\ & + \frac{1}{s(\theta^{[N-1, 1]})} \begin{pmatrix} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T \left(\frac{-\sin \theta_-^{[N-1, 1]}}{c_+^{[N-1, 1]}} \mathbf{q}'_{+^{[N-1, 1]}} + \frac{\sin \theta_+^{[N-1, 1]}}{c_-^{[N-1, 1]}} \mathbf{q}'_{-^{[N-1, 1]}} \right) \\ [W_{\bar{\boldsymbol{\gamma}}'}^{[N-1, 1]}]^T \left(\frac{\cos \theta_-^{[N-1, 1]}}{c_+^{[N-1, 1]}} \mathbf{q}'_{+^{[N-1, 1]}} + \frac{-\cos \theta_+^{[N-1, 1]}}{c_-^{[N-1, 1]}} \mathbf{q}'_{-^{[N-1, 1]}} \right) \end{pmatrix} + \\ & + \begin{pmatrix} \mathbf{0} \\ \frac{1}{c^{[N-2, 2]}} [W_{\bar{\boldsymbol{\gamma}}'}^{[N-2, 2]}]^T \mathbf{q}'^{[N-2, 2]} \end{pmatrix}. \end{aligned} \quad (150)$$

Thus from Eqs. (22), (29), (32), (33) and (150) we find that

$$\mathbf{y} = \begin{pmatrix} \mathbf{r} \\ \boldsymbol{\gamma} \end{pmatrix} = \mathbf{y}_\infty + \frac{1}{\sqrt{D}} \left(\sum_{\alpha=\left\{ \begin{smallmatrix} [N], \\ [N-1, 1] \end{smallmatrix} \right\}} \sum_{\xi} \sum_{\tau=\pm} \tau \mathbf{y}'_{\xi}{}^{\alpha} + \sum_{\xi} \mathbf{y}'_{\xi}{}^{[N-2, 2]} \right) \quad (151)$$

where

$$\mathbf{y}_\infty = \begin{pmatrix} D^2 \bar{a}_{ho} \bar{\mathbf{r}}'_\infty \mathbf{1}_{\bar{\mathbf{r}}'} \\ \bar{\boldsymbol{\gamma}}'_\infty \mathbf{1}_{\bar{\boldsymbol{\gamma}}'} \end{pmatrix}, \quad (152)$$

with $\mathbf{1}_{\bar{\mathbf{r}}'}$ and $\mathbf{1}_{\bar{\boldsymbol{\gamma}}'}$ given in Eqs. (116) and (117) respectively. The vectors $+\mathbf{y}'_{\xi}{}^{\alpha}$, $-\mathbf{y}'_{\xi}{}^{\alpha}$ and $\mathbf{y}'_{\xi}{}^{[N-2, 2]}$ are

$$+\mathbf{y}'_{\xi}{}^{\alpha} = \frac{1}{s(\theta^{\alpha})} \begin{pmatrix} D^2 \bar{a}_{ho} \frac{-\sin \theta_{+}^{\alpha}}{c_{+}^{\alpha}} [\mathbf{q}'_{+}]_{\xi} [(W_{\bar{\mathbf{r}}'}^{\alpha})_{\xi}]^T \\ \frac{\cos \theta_{+}^{\alpha}}{c_{+}^{\alpha}} [\mathbf{q}'_{+}]_{\xi} [(W_{\bar{\boldsymbol{\gamma}}'}^{\alpha})_{\xi}]^T \end{pmatrix}, \quad (153)$$

$$-\mathbf{y}'_{\xi}{}^{\alpha} = \frac{1}{s(\theta^{\alpha})} \begin{pmatrix} D^2 \bar{a}_{ho} \frac{\sin \theta_{+}^{\alpha}}{c_{+}^{\alpha}} [\mathbf{q}'_{-}]_{\xi} [(W_{\bar{\mathbf{r}}'}^{\alpha})_{\xi}]^T \\ -\frac{\cos \theta_{+}^{\alpha}}{c_{+}^{\alpha}} [\mathbf{q}'_{-}]_{\xi} [(W_{\bar{\boldsymbol{\gamma}}'}^{\alpha})_{\xi}]^T \end{pmatrix} \quad (154)$$

for $\alpha = [N]$ or $[N-1, 1]$, and

$$\mathbf{y}'_{\xi}{}^{[N-2, 2]} = \begin{pmatrix} \mathbf{0} \\ \frac{1}{c^{[N-2, 2]}} [\mathbf{q}'^{[N-2, 2]}]_{\xi} [(W_{\bar{\boldsymbol{\gamma}}'}^{[N-2, 2]})_{\xi}]^T \end{pmatrix}. \quad (155)$$

Equations (151), (152), (153), (154) and (155) express, in terms of the internal coordinates $\mathbf{r}$ and $\boldsymbol{\gamma}$, the motion associated with the normal coordinates, $\mathbf{q}'$, about the Lewis structure configuration $\mathbf{y}_{\infty}$.

7 Determining the Symmetry Coordinates

7.1 Primitive Irreducible Coordinates

We determine the symmetry coordinates, and hence the W_r and W_{γ} of Eqs. (86) and (92) respectively, in a two-step process:

- a). Determine two sets of linear combinations of the elements of coordinate vector $\bar{\mathbf{r}}'$ which transform under particular non-orthogonal $[N]$ and $[N-1, 1]$ irreducible representations of S_N . Using these two sets of coordinates we then determine two sets of linear combinations of the elements of coordinate vector $\bar{\boldsymbol{\gamma}}'$ which transform under exactly the same non-orthogonal $[N]$ and $[N-1, 1]$ irreducible representations of S_N as the coordinate sets in the $\bar{\mathbf{r}}'$ sector. In this way we satisfy item 1 of Subsec. 6.2 above. Then another set of linear combinations of the elements of coordinate vector $\bar{\boldsymbol{\gamma}}'$ which transforms under a particular non-orthogonal $[N-2, 2]$ irreducible representation of S_N . These sets of coordinates, which we term primitive irreducible coordinates for reasons discussed below, have W matrices which automatically satisfy Eq. (D4).
- b). Appropriate linear combinations within each coordinate set from item a). above are taken so that the results transform under orthogonal irreducible representations of S_N . These are then the symmetry coordinates of Section 6 above. Care is taken to ensure that the transformation from the coordinates which transform under the non-orthogonal irreducible representations of S_N of item a). above, to the symmetry coordinates which transform from the under orthogonal

irreducible representations of S_N preserve the identity of equivalent representations in the $\bar{\mathbf{r}}'$ and $\bar{\gamma}'$ sectors to ensure that item 1 of Subsec. 6.2 above continues to be satisfied. In this way both items 1 and 2 of Subsec. 6.2 above are satisfied.

What are the advantages of this two-step process? In step a). above we set out to find sets of coordinates transforming irreducibly under S_N which have the simplest functional form possible. This is why we call them primitive irreducible coordinates. In step b). above, we find appropriate linear combinations of the primitive irreducible coordinates so that they transform under orthogonal representations of S_N . Thus the symmetry coordinates are composed of the building blocks of the primitive irreducible coordinates. Furthermore, we choose one of the symmetry coordinates to be just a single primitive irreducible coordinate, and so it describes the simplest motion possible under the requirement that it transforms irreducibly under S_N . The succeeding symmetry coordinate is then chosen to be composed of two primitive invariant coordinates and so on. In this way the complexity of the motions described by the symmetry coordinates is kept to a minimum and only builds up slowly as more symmetry coordinates of a given species are considered.

7.2 The Primitive Irreducible Coordinate Vector, $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}$.

The primitive irreducible coordinates, $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}$, of the $\bar{\mathbf{r}}'$ sector which transform under irreducible, though non-orthogonal, representations of the group S_N are given by

$$\bar{\mathbf{S}}_{\bar{\mathbf{r}}'} = \bar{\mathbf{W}}_{\bar{\mathbf{r}}'} \bar{\mathbf{r}}', \quad (156)$$

where

$$\bar{\mathbf{W}}_{\bar{\mathbf{r}}'} = \begin{pmatrix} \bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N]} \\ \bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N-1, 1]} \end{pmatrix}, \quad (157)$$

$\bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N]}$ is a $1 \times N$ dimensional matrix and $\bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N-1, 1]}$ is an $(N-1) \times N$ dimensional matrix. Hence we can write

$$\bar{\mathbf{S}}_{\bar{\mathbf{r}}'} = \begin{pmatrix} \bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N]} \\ \bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N-1, 1]} \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N]} \bar{\mathbf{r}}' \\ \bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{[N-1, 1]} \bar{\mathbf{r}}' \end{pmatrix}. \quad (158)$$

Since $\bar{\mathbf{W}}_{\bar{\mathbf{r}}'}$ is a matrix which effects the reduction of the representation, $D_{\bar{\mathbf{r}}'}(R)$, under which $\bar{\mathbf{r}}'$ transforms, to particular irreducible, though non-orthogonal, representations of the group S_N under which $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}$ transforms, then

$$\bar{\mathbf{W}}_{\bar{\mathbf{r}}'} D_{\bar{\mathbf{r}}'}(R) \bar{\mathbf{W}}_{\bar{\mathbf{r}}'}^{-1} \bar{\mathbf{S}}_{\bar{\mathbf{r}}'} = \bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}(R) \bar{\mathbf{S}}_{\bar{\mathbf{r}}'}, \quad (159)$$

where

$$\bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}(R) = \left(\begin{array}{c|c} \bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N]}(R) & \mathbf{0} \\ \hline \mathbf{0} & \bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R) \end{array} \right). \quad (160)$$

In the above, $\bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N]}(R)$ is the one-dimensional irreducible matrix representation $[N]$ of S_N and $\bar{\mathbf{D}}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R)$ is an $(N-1) \times (N-1)$ -dimensional, non-orthogonal, irreducible matrix representation $[N-1, 1]$ of S_N .

Thus the primitive irreducible coordinate column vector $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N]}$ is a one-element vector and transforms under $\bar{D}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N]}(R)$, a one-dimensional non-orthogonal irreducible matrix representation $[N]$ of S_N . The symmetry coordinate column vector $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N-1, 1]}$ is a $(N-1)$ -element vector and transforms under $\bar{D}_{\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}}^{[N-1, 1]}(R)$, the aforementioned $(N-1) \times (N-1)$ -dimensional, non-orthogonal, irreducible matrix representation $[N-1, 1]$ of S_N .

7.3 The Primitive Irreducible Coordinate Vector, $\bar{\mathbf{S}}_{\bar{\gamma}'}$.

Likewise the primitive irreducible coordinates, $\bar{\mathbf{S}}_{\bar{\gamma}'}$, of the $\bar{\gamma}'$ sector which transform under irreducible, though non-orthogonal, representations of the group S_N are given by

$$\bar{\mathbf{S}}_{\bar{\gamma}'} = \bar{W}_{\bar{\gamma}'} \bar{\gamma}' \quad (161)$$

and so writing

$$\bar{W}_{\bar{\gamma}'} = \begin{pmatrix} \bar{W}_{\bar{\gamma}'}^{[N]} \\ \bar{W}_{\bar{\gamma}'}^{[N-1, 1]} \\ \bar{W}_{\bar{\gamma}'}^{[N-2, 2]} \end{pmatrix}, \quad (162)$$

where $\bar{W}_{\bar{\gamma}'}^{[N]}$ is a $1 \times N(N-1)/2$ dimensional matrix, $\bar{W}_{\bar{\gamma}'}^{[N-1, 1]}$ is an $(N-1) \times N(N-1)/2$ dimensional matrix and $\bar{W}_{\bar{\gamma}'}^{[N-2, 2]}$ is an $N(N-3)/2 \times N(N-1)/2$ dimensional matrix, then

$$\bar{\mathbf{S}}_{\bar{\gamma}'} = \begin{pmatrix} \bar{\mathbf{S}}_{\bar{\gamma}'}^{[N]} \\ \bar{\mathbf{S}}_{\bar{\gamma}'}^{[N-1, 1]} \\ \bar{\mathbf{S}}_{\bar{\gamma}'}^{[N-2, 2]} \end{pmatrix} = \begin{pmatrix} \bar{W}_{\bar{\gamma}'}^{[N]} \bar{\gamma}' \\ \bar{W}_{\bar{\gamma}'}^{[N-1, 1]} \bar{\gamma}' \\ \bar{W}_{\bar{\gamma}'}^{[N-2, 2]} \bar{\gamma}' \end{pmatrix}, \quad (163)$$

where

$$\bar{W}_{\bar{\gamma}'}^\alpha \bar{\gamma}' = \sum_{j=1}^N \sum_{i < j} [\bar{W}_{\bar{\gamma}'}^\alpha]_{ij} \bar{\gamma}'_{ij}. \quad (164)$$

Since $\bar{W}_{\bar{\gamma}'}$ is a matrix which effects the reduction of the representation, $\bar{D}_{\bar{\gamma}'}(R)$, under which the $\bar{\gamma}'$ vector transforms, to particular irreducible, though non-orthogonal, representations of the group S_N under which $\bar{\mathbf{S}}_{\bar{\gamma}'}$ transforms, then

$$\bar{W}_{\bar{\gamma}'} D_{\bar{\gamma}'}(R) \bar{W}_{\bar{\gamma}'}^{-1} \bar{\mathbf{S}}_{\bar{\gamma}'} = \bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}(R) \bar{\mathbf{S}}_{\bar{\gamma}'}, \quad (165)$$

where

$$\bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}(R) = \left(\begin{array}{c|cc} \bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N]}(R) & \mathbf{0} & \mathbf{0} \\ \hline \mathbf{0} & \bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N-1, 1]}(R) & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & \bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N-2, 2]}(R) \end{array} \right). \quad (166)$$

In the above, $\bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N]}(R)$ is the one-dimensional irreducible matrix representation $[N]$ of S_N , $\bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N-1, 1]}(R)$ is an $(N-1) \times (N-1)$ -dimensional, non-orthogonal, irreducible matrix representation $[N-1, 1]$ of S_N and $\bar{D}_{\bar{\mathbf{S}}_{\bar{\gamma}'}}^{[N-2, 2]}(R)$ is an $\{N(N-3)/2\} \times \{N(N-3)/2\}$ -dimensional, non-orthogonal, irreducible matrix representation $[N-2, 2]$ of S_N .

Note that according to item a). above,

$$\overline{D}_{\overline{S}_{\overline{\tau}'}}^{\alpha}(R) = \overline{D}_{\overline{S}_{\overline{\tau}'}}^{\alpha}(R) = \overline{D}^{\alpha}(R), \quad (167)$$

where α is the partition $[N]$ or $[N-1, 1]$.

Thus primitive irreducible coordinate column vector $\overline{S}_{\overline{\tau}'}^{[N]}$ is a one-element vector and transforms under $\overline{D}^{[N]}(R)$, the one-dimensional irreducible matrix representation $[N]$ of S_N , while the primitive irreducible coordinate column vector $\overline{S}_{\overline{\tau}'}^{[N-1, 1]}$ is an $(N-1)$ -element vector and transforms under $\overline{D}^{[N-1, 1]}(R)$, an $(N-1) \times (N-1)$ -dimensional, non-orthogonal, irreducible matrix representation $[N-1, 1]$ of S_N . The primitive irreducible coordinate column vector $\overline{S}_{\overline{\tau}'}^{[N-2, 2]}$ is an $N(N-3)/2$ -element vector and transforms under $\overline{D}^{[N-2, 2]}(R)$, the $\{N(N-3)/2\} \times \{N(N-3)/2\}$ -dimensional, non-orthogonal, irreducible matrix representation $[N-2, 2]$ of S_N .

We also note that according to Eq. (D4)

$$\overline{W}_{\mathbf{X}'}^{\alpha} [\overline{W}_{\mathbf{X}'}^{\alpha'}]^T = 0, \quad \forall \quad \alpha \neq \alpha'. \quad (168)$$

7.4 The Full Primitive Irreducible Coordinate Vector, $\overline{S}$.

Like for the discussion of the symmetry coordinates in Subsection 6.1 above, it is useful to form a full primitive irreducible coordinate vector that groups primitive irreducible coordinates of the same species together as follows:

$$\overline{S} = P \begin{pmatrix} \overline{S}_{\overline{\tau}'} \\ \overline{S}_{\overline{\tau}'} \end{pmatrix} = \begin{pmatrix} \overline{S}_{\overline{\tau}'}^{[N]} \\ \overline{S}_{\overline{\tau}'}^{[N-1, 1]} \\ \overline{S}_{\overline{\tau}'}^{[N-1, 1]} \\ \overline{S}_{\overline{\tau}'}^{[N-2, 2]} \end{pmatrix} = \begin{pmatrix} \overline{S}^{[N]} \\ \overline{S}^{[N-1, 1]} \\ \overline{S}^{[N-2, 2]} \end{pmatrix}, \quad (169)$$

where the orthogonal matrix P is given by Eq. (97) and

$$\overline{S}^{[N]} = \begin{pmatrix} \overline{S}_{\overline{\tau}'}^{[N]} \\ \overline{S}_{\overline{\tau}'}^{[N]} \end{pmatrix}, \quad \overline{S}^{[N-1, 1]} = \begin{pmatrix} \overline{S}_{\overline{\tau}'}^{[N-1, 1]} \\ \overline{S}_{\overline{\tau}'}^{[N-1, 1]} \end{pmatrix} \quad \text{and} \quad \overline{S}^{[N-2, 2]} = \overline{S}_{\overline{\tau}'}^{[N-2, 2]}. \quad (170)$$

Now consider applying a transformation $\overline{W}$ to Eqs. (46), (47), (48) and (49), where

$$\overline{W} = P \left(\overline{W}_{\overline{\tau}'} \middle| \begin{array}{c} \mathbf{0} \\ \overline{W}_{\overline{\tau}'} \end{array} \right) = \begin{pmatrix} \overline{W}_{\overline{\tau}'}^{[N]} & \mathbf{0} \\ \mathbf{0} & \overline{W}_{\overline{\tau}'}^{[N]} \\ \overline{W}_{\overline{\tau}'}^{[N-1, 1]} & \mathbf{0} \\ \mathbf{0} & \overline{W}_{\overline{\tau}'}^{[N-1, 1]} \\ \mathbf{0} & \overline{W}_{\overline{\tau}'}^{[N-2, 2]} \end{pmatrix}, \quad (171)$$

where

$$\overline{S} = \overline{W} \mathbf{y}'. \quad (172)$$

Eq. (172) implies that the representation under which the displacement coordinate vector $\mathbf{S}$ transforms is

$$\overline{D}_{\overline{S}} = \overline{W} D_{\mathbf{y}'}(R) \overline{W}^{-1}, \quad (173)$$

where $\overline{D}_{\mathbf{y}'}(R)$ is given by Eq. (E1). The non-orthogonal matrix representation $D_{\overline{S}}(R)$ is a block diagonal matrix of irreducible, though non-orthogonal, representations of the form

$$\begin{aligned} \overline{D}_{\overline{S}}(R) &= P \left(\begin{array}{c|c} \overline{D}_{\overline{S}_{\mathbf{r}'}}(R) & \mathbf{0} \\ \hline \mathbf{0} & \overline{D}_{\overline{S}_{\overline{\gamma}'}}(R) \end{array} \right) P^T \\ &= \left(\begin{array}{cc|cc|c} \overline{D}^{[N]}(R) & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \overline{D}^{[N]}(R) & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & \overline{D}^{[N-1, 1]}(R) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \overline{D}^{[N-1, 1]}(R) & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \overline{D}^{[N-2, 2]}(R) \end{array} \right), \end{aligned} \quad (174)$$

where we have also used Eq. (167) from above in the last step.

7.5 The Transformation, U , from Primitive Irreducible Coordinates to Symmetry Coordinates.

The symmetry coordinate vector $\mathbf{S}$ will be related to the primitive irreducible coordinate vector $\overline{\mathbf{S}}$ by a non-orthogonal linear transformation U , i.e.

$$\mathbf{S} = U \overline{\mathbf{S}} \quad (175)$$

and likewise $D_S(R)$ will be related to $\overline{D}_{\overline{S}}(R)$ by

$$D_S(R) = U \overline{D}_{\overline{S}}(R) U^{-1}. \quad (176)$$

Thus from Eqs. (100), (172), (E2) and (173), W will be related to $\overline{W}$ by the same transformation, i.e.

$$W = U \overline{W}, \quad (177)$$

where

$$U = \left(\begin{array}{cc|cc|c} U_{\overline{\mathbf{r}'}}^{[N]} & 0 & 0 & 0 & 0 \\ 0 & U_{\overline{\gamma}'}^{[N]} & 0 & 0 & 0 \\ \hline 0 & 0 & U_{\overline{\mathbf{r}'}}^{[N-1, 1]} & 0 & 0 \\ 0 & 0 & 0 & U_{\overline{\gamma}'}^{[N-1, 1]} & 0 \\ \hline 0 & 0 & 0 & 0 & U_{\overline{\gamma}'}^{[N-2, 2]} \end{array} \right), \quad (178)$$

so that

$$W = \left(\begin{array}{cc|cc} U_{\overline{\mathbf{r}'}}^{[N]} \overline{W}_{\overline{\mathbf{r}'}}^{[N]} & \mathbf{0} & & \\ \mathbf{0} & U_{\overline{\gamma}'}^{[N]} \overline{W}_{\overline{\gamma}'}^{[N]} & & \\ \hline U_{\overline{\mathbf{r}'}}^{[N-1, 1]} \overline{W}_{\overline{\mathbf{r}'}}^{[N-1, 1]} & \mathbf{0} & & \\ \mathbf{0} & U_{\overline{\gamma}'}^{[N-1, 1]} \overline{W}_{\overline{\gamma}'}^{[N-1, 1]} & & \\ \hline \mathbf{0} & U_{\overline{\gamma}'}^{[N-2, 2]} \overline{W}_{\overline{\gamma}'}^{[N-2, 2]} & & \end{array} \right). \quad (179)$$

Note that from Eqs. (E8), (174) and (176),

$$U_{\vec{\gamma}'}^\alpha = A_U^\alpha U_{\vec{r}'}^\alpha, \quad (180)$$

where α is $[N]$ or $[N-1, 1]$ and A_U^α is a number (see below).

Comparing Eq. (179) with Eq. (99) we see that

$$W_{\mathbf{X}'}^\alpha = U_{\mathbf{X}'}^\alpha \overline{W}_{\mathbf{X}'}^\alpha, \quad (181)$$

and according to Eq. (102) $U_{\mathbf{X}'}^\alpha$ must satisfy

$$U_{\mathbf{X}'}^\alpha \left\{ \overline{W}_{\mathbf{X}'}^\alpha [\overline{W}_{\mathbf{X}'}^\alpha]^T \right\} [U_{\mathbf{X}'}^\alpha]^T = I_\alpha. \quad (182)$$

Thus to determine $U_{\mathbf{X}'}^\alpha$, we need to know $\overline{W}_{\mathbf{X}'}^\alpha [\overline{W}_{\mathbf{X}'}^\alpha]^T$ beforehand. Note that when we use Eq. (180) in Eq. (182) we must have

$$\overline{W}_{\vec{\gamma}'}^\alpha [\overline{W}_{\vec{\gamma}'}^\alpha]^T = \frac{1}{(A_U^\alpha)^2} \overline{W}_{\vec{r}'}^\alpha [\overline{W}_{\vec{r}'}^\alpha]^T \quad (183)$$

when α is $[N]$ or $[N-1, 1]$.

7.6 Symmetry Coordinates Belonging to the $[N]$ Species

7.6.1 The $\vec{r}'$ Sector

The Primitive Irreducible Coordinate The primitive irreducible coordinate $\overline{\mathbf{S}}_{\vec{r}'}^{[N]}$ which is invariant under S_N is readily seen to be

$$\overline{\mathbf{S}}_{\vec{r}'}^{[N]} = \sum_{i=1}^N \vec{r}'_i. \quad (184)$$

From Eq. (158) we can identify $\overline{W}_{\vec{r}'}^{[N]}$ as

$$[\overline{W}_{\vec{r}'}^{[N]}]_i = [\mathbf{1}_{\vec{r}'}]_i \quad (185)$$

where $\mathbf{1}_{\vec{r}'}$ is defined in Eq. (116). We leave the calculation of $\overline{\mathbf{S}}_{\vec{r}'}^{[N-1, 1]}$ and $\overline{W}_{\vec{r}'}^{[N-1, 1]}$ to a subsequent paper[10].

The Symmetry Coordinate One readily sees that

$$\overline{W}_{\vec{r}'}^{[N]} [\overline{W}_{\vec{r}'}^{[N]}]^T = N \quad (186)$$

and so from Eq. (182)

$$U_{\vec{r}'}^{[N]} = \frac{1}{\sqrt{N}}. \quad (187)$$

Thus from Eqs. (175), (181), (184), (185) and (187)

$$[W_{\vec{r}'}^{[N]}]_i = \frac{1}{\sqrt{N}} [\mathbf{1}_{\vec{r}'}]_i \quad (188)$$

and

$$\mathbf{S}_{\vec{r}'}^{[N]} = \frac{1}{\sqrt{N}} \sum_{i=1}^N \vec{r}'_i. \quad (189)$$

We leave the calculation of $W_{\vec{r}'}^{[N-1, 1]}$ and $\mathbf{S}_{\vec{r}'}^{[N-1, 1]}$ to a subsequent paper[10].

Motions Associated with Symmetry Coordinate $\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}$. According to Eqs. (113), (116), (114) and (188) the motions associated with symmetry coordinate $\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}$ in the unscaled internal displacement coordinates $\mathbf{r}$ about the unscaled Lewis structure configuration

$$\mathbf{r}_\infty = D^2 \bar{a}_{ho} \bar{\mathbf{r}}'_\infty \mathbf{1}_{\bar{\mathbf{r}}'} \quad (190)$$

are given by

$$\mathbf{r}^{[N]} = \bar{a}_{ho} D^{3/2} \bar{\mathbf{r}}'^{[N]} = \bar{a}_{ho} D^{3/2} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]} [(W_{\bar{\mathbf{r}}'}^{[N]})]^T = \bar{a}_{ho} \sqrt{\frac{D^3}{N}} \mathbf{S}_{\bar{\mathbf{r}}'}^{[N]} \mathbf{1}_{\bar{\mathbf{r}}'}. \quad (191)$$

Thus we see that the motions associated with symmetry coordinate $\mathbf{S}_{\bar{\mathbf{r}}'}^{[N]}$ involve symmetric-stretch motions about the Lewis structure configuration, where all of the radii expand and contract together.

We leave the calculation of $\mathbf{r}^{[N-1, 1]}$ to a subsequent paper[10].

7.6.2 The $\bar{\gamma}'$ Sector

The Primitive Irreducible Coordinate According to item a). above, $\bar{\mathbf{S}}_{\bar{\gamma}'}^{[N]}$ should transform under exactly the same non-orthogonal irreducible $[N]$ representation of S_N as $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N]}$. Since $\bar{\mathbf{S}}_{\bar{\mathbf{r}}'}^{[N]}$ is invariant under all of the elements, we simply require that $\bar{\mathbf{S}}_{\bar{\gamma}'}^{[N]}$ be invariant under S_N .

Now according to Eq. (4)

$$\gamma_{ij} = \hat{\mathbf{r}}_i \cdot \hat{\mathbf{r}}_j, \quad (192)$$

where $\hat{\mathbf{r}}_i$ is the unit vector from the center of the confining field to particle i . Now $\sum_{i=1}^N \hat{\mathbf{r}}_i$ is invariant under S_N and so $(\sum_{i=1}^N \hat{\mathbf{r}}_i) \cdot (\sum_{j=1}^N \hat{\mathbf{r}}_j)$ is invariant under S_N . The contribution from the terms with $i = j$, i.e. $\sum_{i=1}^N \hat{\mathbf{r}}_i \cdot \hat{\mathbf{r}}_i = N$, is also invariant on its own. Since this is just equal to the number N , we subtract this term from $(\sum_{i=1}^N \hat{\mathbf{r}}_i) \cdot (\sum_{j=1}^N \hat{\mathbf{r}}_j)$ and consider the invariant quantity $(\sum_{i=1}^N \hat{\mathbf{r}}_i) \cdot (\sum_{j \neq i}^N \hat{\mathbf{r}}_j)$. According to Eq. (192) this equals $\sum_{i=1}^N \sum_{j \neq i} \gamma_{ij} = 2 \sum_{j=2}^N \sum_{i < j} \gamma_{ij}$ and upon using Eq. (33) we obtain $[N(N-1)/2] \bar{\gamma}_\infty + \delta^{1/2} \sum_{j=2}^N \sum_{i < j} \bar{\gamma}'_{ij}$. Since the first term is invariant under S_N , $\sum_{j=2}^N \sum_{i < j} \bar{\gamma}'_{ij}$ is invariant under S_N , and so we identify the primitive irreducible coordinate of the $\bar{\gamma}'$ sector which transforms under the $[N]$ representation as

$$\bar{\mathbf{S}}_{\bar{\gamma}'}^{[N]} = \sum_{j=2}^N \sum_{i < j} \bar{\gamma}'_{ij}. \quad (193)$$

From Eqs. (40), (163), (164) and (193)

$$[\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N]}]_{ij} = [\mathbf{1}_{\bar{\gamma}'}]_{ij}, \quad (194)$$

where $\mathbf{1}_{\bar{\gamma}'}$ is defined in Eq. (117). We leave the calculation of $\bar{\mathbf{S}}_{\bar{\gamma}'}^{[N-1, 1]}$, $\bar{\mathbf{S}}_{\bar{\gamma}'}^{[N-2, 2]}$, $\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N-1, 1]}$ and $\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N-2, 2]}$ to a subsequent paper[10].

The Symmetry Coordinate One readily sees that

$$\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N]} [\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N]}]^T = \sum_{j=1}^N \sum_{i < j} [\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N]}]_{ij} [\bar{\mathbf{W}}_{\bar{\gamma}'}^{[N]}]_{ij} = \frac{N(N-1)}{2}, \quad (195)$$

where we have used Eqs. (194) and (117) in the last step, and so from Eq. (182)

$$U_{\vec{\gamma}'}^{[N]} = \sqrt{\frac{2}{N(N-1)}}. \quad (196)$$

Thus from Eqs. (175), (181), (193), (194) and (196)

$$[W_{\vec{\gamma}'}^{[N]}]_{ij} = \sqrt{\frac{2}{N(N-1)}} [\mathbf{1}_{\vec{\gamma}'}]_{ij} \quad (197)$$

and

$$\mathbf{S}_{\vec{\gamma}'}^{[N]} = \sqrt{\frac{2}{N(N-1)}} \sum_{j=2}^N \sum_{i < j} \vec{\gamma}'_{ij}. \quad (198)$$

We leave the calculation of $W_{\vec{\gamma}'}^{[N-1, 1]}$, $W_{\vec{\gamma}'}^{[N-2, 2]}$, $\mathbf{S}_{\vec{\gamma}'}^{[N-1, 1]}$ and $\mathbf{S}_{\vec{\gamma}'}^{[N-2, 2]}$ to a subsequent paper[10].

Motions Associated with Symmetry Coordinate $\mathbf{S}_{\vec{\gamma}'}^{[N]}$. According to Eqs. (113), (115), (117) and (197) the motions associated with symmetry coordinate $\mathbf{S}_{\vec{\gamma}'}^{[N]}$ in the unscaled internal displacement coordinates γ about the Lewis structure configuration

$$\gamma_{\infty} = \bar{\gamma}_{\infty} \mathbf{1}_{\vec{\gamma}'} \quad (199)$$

are given by

$$\gamma^{[N]} = \frac{1}{\sqrt{D}} \gamma'^{[N]} = \frac{1}{\sqrt{D}} \mathbf{S}_{\vec{\gamma}'}^{[N]} [(W_{\vec{\gamma}'}^{[N]})]^T = \sqrt{\frac{2}{N(N-1)D}} \mathbf{S}_{\vec{\gamma}'}^{[N]} \mathbf{1}_{\vec{\gamma}'}. \quad (200)$$

Thus we see that the motions associated with symmetry coordinate $\mathbf{S}_{\vec{\gamma}'}^{[N]}$ involve symmetric-bend motions about the Lewis structure configuration, where all of the angles open out and contract together like a parasol, but with the radii unchanged.

We leave the calculation of $\gamma^{[N-1, 1]}$ and $\gamma^{[N-2, 2]}$ to a subsequent paper[10].

8 The Frequencies and Normal-Mode Coordinates of the [N] Species of the System.

8.1 The G and FG matrices in the Symmetry-Coordinate Basis.

We can use the $W_{\vec{\gamma}'}^{[N]}$ matrices of Eqs. (188) and (197) to calculate the reduced $\mathbf{G}$ and $\mathbf{FG}$ matrix elements, $[\sigma_{[N]}^{\mathbf{G}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$ and $[\sigma_{[N]}^{\mathbf{FG}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$, of Eq. (120). An outline of this calculation is to be found in Appendix F, with the following results.

8.1.1 The Matrix Elements $[\sigma_{[N]}^{\mathbf{G}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$.

Using Eqs. (59), (60), (61), (62), (63), (66), (67), (68), (120), (188) and (197) we derive

$$\sigma_{[N]}^{\mathbf{G}} = \begin{pmatrix} [\sigma_{[N]}^{\mathbf{G}}]_{\vec{r}', \vec{r}'} = \tilde{a}' & [\sigma_{[N]}^{\mathbf{G}}]_{\vec{r}', \vec{\gamma}'} = 0 \\ [\sigma_{[N]}^{\mathbf{G}}]_{\vec{\gamma}', \vec{r}'} = 0 & [\sigma_{[N]}^{\mathbf{G}}]_{\vec{\gamma}', \vec{\gamma}'} = (\tilde{g}' + 2(N-1)\tilde{h}') \end{pmatrix}. \quad (201)$$

We leave the calculation of the matrix elements $[\sigma_{[N-1, 1]}^G]_{\mathbf{x}'_1, \mathbf{x}'_2}$ and $\sigma_{[N-2, 2]}^G = [\sigma_{[N-2, 2]}^{FG}]_{\bar{\gamma}', \bar{\gamma}'}$ to a subsequent paper[10].

8.1.2 The Matrix Elements $[\sigma_{[N]}^{FG}]_{\mathbf{x}'_1, \mathbf{x}'_2}$

Using Eqs. (59), (60), (61), (62), (63), (64), (65), (120), (135), (188) and (197) we derive

$$\sigma_{[N]}^{FG} = \begin{pmatrix} [\sigma_{[N]}^{FG}]_{\bar{r}', \bar{r}'} = (\tilde{a} + \tilde{b}N) & [\sigma_{[N]}^{FG}]_{\bar{r}', \bar{\gamma}'} = \sqrt{2(N-1)} \left(\tilde{e} + \frac{N}{2} \tilde{f} \right) \\ [\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{r}'} = \sqrt{2(N-1)} \left(\tilde{c} + \frac{N}{2} \tilde{d} \right) & [\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} = \left(\tilde{g} + 2(N-1)\tilde{h} + \frac{N(N-1)}{2} \tilde{i} \right) \end{pmatrix}. \quad (202)$$

We leave the calculation of the matrix elements $[\sigma_{[N-1, 1]}^{FG}]_{\mathbf{x}'_1, \mathbf{x}'_2}$ and $\sigma_{[N-2, 2]}^{FG} = [\sigma_{[N-2, 2]}^{FG}]_{\bar{\gamma}', \bar{\gamma}'}$ to a subsequent paper[10].

8.2 The Frequencies and Normal Modes.

Using Eqs. (202) in Eq. (137) we obtain $\lambda_{[N]}^{\pm}$. The frequencies are then determined from Eq. (52). The $\bar{r}'$ - $\bar{\gamma}'$ mixing angles, $\theta_{\pm}^{[N]}$ for the $[N]$ species are determined from Eq. (138). The normalization constant $c^{[N]}$ of the reduced normal-coordinate coefficient vector, $\mathbf{c}^{[N]}$, of Eqs. (124) and (129) are determined from Eqs. (141) and (142). One then determines the normal mode vector, $\mathbf{q}'$, through Eqs. (145), (189) and (198). Thus we arrive at

$$q_+^{[N]} = c_+^{[N]} \cos \theta_+^{[N]} \frac{1}{\sqrt{N}} \sum_{i=1}^N \bar{r}'_i + c_+^{[N]} \sin \theta_+^{[N]} \sqrt{\frac{2}{N(N-1)}} \sum_{j=2}^N \sum_{i < j} \bar{\gamma}'_{ij} \quad (203)$$

$$q_-^{[N]} = c_-^{[N]} \cos \theta_-^{[N]} \frac{1}{\sqrt{N}} \sum_{i=1}^N \bar{r}'_i + c_-^{[N]} \sin \theta_-^{[N]} \sqrt{\frac{2}{N(N-1)}} \sum_{j=2}^N \sum_{i < j} \bar{\gamma}'_{ij} \quad (204)$$

We leave the calculation of $\mathbf{q}'_+^{[N-1, 1]}$, $\mathbf{q}'_-^{[N-1, 1]}$ and $\mathbf{q}'^{[N-2, 2]}$ to a subsequent paper[10].

8.3 The Motions Associated with the Normal Modes.

From Eqs. (151), (152), (153), (154), (155), (188) and (197)

$$\mathbf{y} = \begin{pmatrix} \mathbf{r} \\ \gamma \end{pmatrix} = \mathbf{y}_{\infty} + \sum_{\alpha \in \left\{ \begin{smallmatrix} [N], \\ [N-1, 1] \end{smallmatrix} \right\}} \sum_{\xi} \sum_{\tau=\pm} \begin{pmatrix} \tau \mathbf{r}_{\xi}^{\alpha} \\ \tau \gamma_{\xi}^{\alpha} \end{pmatrix} + \sum_{\xi} \begin{pmatrix} \mathbf{0} \\ \gamma_{\xi}^{[N-2, 2]} \end{pmatrix}, \quad (205)$$

where $\mathbf{y}_{\infty}$ is given by Eq. (152). The ξ sum for the $[N]$ species only includes one term and so

$$+\mathbf{r}^{[N]} = -\bar{a}_{ho} \sqrt{\frac{D^3}{N}} \frac{\sin \theta_-^{[N]}}{s(\theta^{[N]}) c_+^{[N]}} q_+^{[N]} \mathbf{1}_{\bar{r}'}, \quad (206)$$

$$+\gamma^{[N]} = \sqrt{\frac{2}{N(N-1)D}} \frac{\cos \theta_-^{[N]}}{s(\theta^{[N]}) c_+^{[N]}} q_+^{[N]} \mathbf{1}_{\bar{\gamma}'}, \quad (207)$$

$$_-\mathbf{r}^{[N]} = \bar{a}_{ho} \sqrt{\frac{D^3}{N}} \frac{\sin \theta_+^{[N]}}{s(\theta^{[N]}) c_-^{[N]}} q_-'^{[N]} \mathbf{1}_{\bar{\mathbf{r}}'}, \quad (208)$$

$$_-\gamma^{[N]} = -\sqrt{\frac{2}{N(N-1)D}} \frac{\cos \theta_+^{[N]}}{s(\theta^{[N]}) c_-^{[N]}} q_-'^{[N]} \mathbf{1}_{\bar{\gamma}'}. \quad (209)$$

We leave the calculation of $_\tau \mathbf{r}_\xi^{[N-1, 1]}$, $_\tau \gamma_\xi^{[N-1, 1]}$ and $\gamma_\xi^{[N-2, 2]}$ to a subsequent paper[10].

9 Summary and Conclusions

In earlier papers[2, 4, 8] we have developed the method of dimensional perturbation theory[3] at low orders to examine the energies of quantum-confined systems such as atoms, quantum dots and Bose-Einstein condensates, both the ground and excited states. Dimensional perturbation theory (DPT) has many advantages. These include the fact that the number of particles, N , enters into the theory as a parameter, and so it is easy to calculate results for an arbitrary number of particles. Also the theory is a beyond-mean-field method, directly accounting for each interaction, rather than some average representation of the interactions. This means that it is appropriate for confined systems of both weakly interacting and strongly interacting particles. In the case of a trapped gaseous atomic Bose-Einstein condensate (BEC) the system is typically a weakly interacting system for which a mean-field approximation is valid. However, if N is increased, or Feshbach resonances are exploited to increase the effective scattering length, or the system is squeezed in one or more directions, it will transition into a strongly-interacting regime where the mean-field approach breaks down. Such systems have been created in the laboratory. Dimensional perturbation theory, however, is equally applicable to both regimes and may be used to study the transition from weak to strongly interacting regimes. Even at lowest order DPT includes beyond-mean-field effects.

In this paper we have begun to address the problem of calculating the zeroth-order DPT wave function for quantum-confined systems, and it is a considerably expanded discussion of work briefly outlined in a previous letter[9]. At large dimensions the Jacobian-weighted wave function of the system becomes harmonic, with the system oscillating about a configuration termed the Lewis structure. Notwithstanding the relatively simple form of the large-dimension, zeroth-order wave function, it includes beyond-mean-field effects.

With the wave function many more properties of the system become accessible. The normal mode coordinates of the zeroth-order wave function reveals the nature of the excitations of the system and the lowest-order wave function yields expectation values and transition matrix elements to zeroth order in DPT. For macroscopic quantum-confined systems, such as the BEC, the wave function is uncloaked in an explicit way as the density profile may be viewed in a direct fashion. This is readily calculable at zeroth order in DPT from the zeroth-order wave function. Also, calculating energies and wave functions to higher orders in $1/D$ requires as input the lowest-order wave function.

In this paper we limit ourselves to setting up the general theory of the zeroth order DPT wave function for a quantum-confined system. We illustrate this by applying it to the breathing and center of mass modes

of the $[N]$ species. In the case of the BEC the ground state energies and excitation frequencies of these modes have been calculated using DPT in a previous paper. The modes of the $[N - 1, 1]$ and $[N - 2, 2]$ species are derived in a subsequent paper. The presence of these additional normal-mode coordinates in the theory indicates the existence of modes of excitation of the BEC beyond those which are typically considered. The zeroth-order density profile of the ground state condensate is to be considered in another paper, and is relatively simple to calculate once the zeroth-order wave function is known.

This paper restricts itself to consideration of quantum-confined systems with a confining potential of spherical symmetry. The extension of this method to systems with cylindrical symmetry is relatively straightforward and is to be discussed in subsequent papers. This is particularly important for the BEC as almost all current laboratory traps have cylindrical symmetry.

As we have discussed in an earlier paper[4] considering energies and excitation frequencies of the BEC, low-order DPT calculations of confined interacting particles, and this includes strongly interacting systems, are accurate out to N equals a few thousand particles. To obtain more accurate energies, and to extend the calculation beyond a few thousand particles we need to go beyond these low-order calculations of the energies and excitation frequencies. These calculations are facilitated by the present paper since, as we have already mentioned, it requires the zeroth-order DPT wave function. This will be pursued in later papers.

10 Acknowledgments

We would like to thank the Army Research Office for ongoing support. This work was also supported in part by the Office of Naval Research.

A The Wilson FG matrix method

In this appendix, we derive the Wilson FG matrix method[14] which is at the heart of our obtaining the normal mode frequencies, the first-order energy correction, the normal coordinates and the wave function at lowest order. The derivation involves a transformation to the set of coordinates called normal-coordinates in which both the differential term and the potential term of Eq. (45) are diagonal. We begin by defining a symmetric transformation, $\mathbf{A}$, that transforms from the vector $\bar{\mathbf{y}}'$, defined by Eq. (38), to $\bar{\mathbf{z}}'$ ($\bar{\mathbf{z}}' = \mathbf{A}\bar{\mathbf{y}}'$). $\mathbf{A}$ is an active transformation and it has the property that it diagonalizes the symmetric $\mathbf{G}$ matrix to unity. That is, $\mathbf{A}$ satisfies

$$\mathbf{A}^T \mathbf{G} \mathbf{A} = \mathbf{I}, \tag{A1}$$

which can also be written as

$$\mathbf{G} = \mathbf{A}^{-1}(\mathbf{A}^{-1})^T, \tag{A2}$$

where we have used the property that $\mathbf{A}$ is symmetric ($\mathbf{A} = \mathbf{A}^T$) in the derivation of Eq. (A2). $\hat{H}_1$ then becomes

$$\hat{H}_1 \rightarrow -\frac{1}{2}\partial_{\bar{\mathbf{z}}'}^T \partial_{\bar{\mathbf{z}}'} + \frac{1}{2}\bar{\mathbf{z}}'^T (\mathbf{A}^{-1})^T \mathbf{F} \mathbf{A}^{-1} \bar{\mathbf{z}}'. \quad (\text{A3})$$

Next we focus our attention on the potential term, the term involving the matrix $\mathbf{F}$. We introduce another transformation, $\mathbf{U}$ ($\mathbf{q}' = \mathbf{U}\bar{\mathbf{z}}'$), that diagonalizes the potential term while simultaneously leaving the differential term unchanged. This orthogonal transformation ($\mathbf{U}^T \mathbf{U} = \mathbf{I}$, where $\mathbf{I}$ is the identity matrix) leaves the differential term in the same form as in Eq. (A3), and the potential term becomes

$$\mathbf{U}(\mathbf{A}^{-1})^T \mathbf{F} \mathbf{A}^{-1} \mathbf{U}^T = \mathbf{\Lambda}, \quad (\text{A4})$$

where $\mathbf{\Lambda}$ is a diagonal matrix. That is,

$$\hat{H}_1 \rightarrow -\frac{1}{2}\partial_{\mathbf{q}'}^T \partial_{\mathbf{q}'} + \frac{1}{2}\mathbf{q}'^T \mathbf{\Lambda} \mathbf{q}'. \quad (\text{A5})$$

From Eq. (A5) it is clear that $\mathbf{\Lambda}$ is the diagonal matrix of the squares of the frequencies of the normal modes (see Eq. (52)).

Now if we introduce the projection operators

$$P_b = \mathbf{b}'' \mathbf{b}''^T, \quad (\text{A6})$$

where the c^{th} element of the column vector $\mathbf{b}''$ is

$$(\mathbf{b}'')_c = \delta_{bc} \quad (\text{A7})$$

with b and c running from 1 to $(N+1)/2$, i.e. all of the elements of the column vector are zero except the b^{th} element which is equal to unity, then

$$\mathbf{\Lambda} \mathbf{b}'' \mathbf{b}''^T \mathbf{q}' = [\mathbf{\Lambda}]_{bb} \mathbf{b}'' [\mathbf{q}']_b = \lambda_b \mathbf{b}'' [\mathbf{q}']_b, \quad (\text{A8})$$

where $[\mathbf{q}']_b$, the b^{th} normal-mode coordinate, is given by

$$[\mathbf{q}']_b = \mathbf{b}''^T \mathbf{q}' \quad (\text{A9})$$

Since $[\mathbf{q}']_b$ is just a multiplier on all sides of Eq. (A8), Eq. (A8) reduces to

$$\mathbf{\Lambda} \mathbf{b}'' = \lambda_b \mathbf{b}'', \quad (\text{A10})$$

Using Eq. (A4), Eq. (A10) reads

$$\mathbf{U}(\mathbf{A}^{-1})^T \mathbf{F} \mathbf{A}^{-1} \mathbf{U}^T \mathbf{b}'' = \lambda_b \mathbf{b}''. \quad (\text{A11})$$

There are two important ways we can proceed from here.

1. If we now define

$$\mathbf{b}'' = \mathbf{U} \mathbf{A} \mathbf{B} \quad (\text{A12})$$

then Eq. (A9), the equation for the b^{th} normal-mode coordinate, reads

$$[\mathbf{q}']_b = \mathcal{B}^T \mathbf{A}^2 \bar{\mathbf{y}}' = \mathcal{B}^T \mathbf{G}^{-1} \bar{\mathbf{y}}', \quad (\text{A13})$$

where we have used Eqs. (A2) and (A4), and from Eq. (A11) we find that $\mathcal{B}$ satisfies the eigenvalue equation

$$\mathbf{G} \mathbf{F} \mathcal{B} = \lambda_b \mathcal{B}. \quad (\text{A14})$$

The normalization of $\mathcal{B}$ is determined by the normalization condition for $\mathbf{b}''$,

$$\mathbf{b}''^T \mathbf{b}'' = 1 \quad (\text{A15})$$

(see Eq. (A7)). Using Eqs. (A2) and (A4) in Eq. (A15) we obtain

$$\mathcal{B}^T \mathbf{G}^{-1} \mathcal{B} = 1. \quad (\text{A16})$$

Notice how the constant matrix $\mathbf{G}^{-1}$ is the contravariant metric tensor for the space. This is to be expected from the form of Eq. (45).

2. All of the work applying DPT to large- N quantum-confined systems up till now has been focussed on calculating energies and adopts the procedure outlined in item 1. The frequency squared, λ_b , is calculated from Eq. (A14) by solving

$$\det(\mathbf{G} \mathbf{F} - \lambda \mathbf{B}) = 0, \quad (\text{A17})$$

Since we are also interested in large- D wave functions and normal coordinates, our life would be easier if $\mathcal{B}$ were the coefficient vector of the displacement coordinates contributing to the normal coordinate $[\mathbf{q}']_b$. Looking at Eq. (A13) we see that this is not so as there is a $\mathbf{G}^{-1}$ operation involved. Thus instead of following the procedure outlined in item 1 above, in this paper we solve directly for the coefficient vector, $\mathbf{b}$, of the displacement coordinates contributing to the normal coordinate $[\mathbf{q}']_b$,

$$\mathbf{b} = \mathbf{G}^{-1} \mathcal{B}. \quad (\text{A18})$$

Thus in terms of $\mathbf{b}$, Eq. (A12) for the normal mode coordinate takes on the desired form

$$[\mathbf{q}']_b = \mathbf{b}^T \bar{\mathbf{y}}'. \quad (\text{A19})$$

Using Eq. (A18) in Eq. (A14) we find that the coefficient vector satisfies the eigenvalue equation

$$\mathbf{F} \mathbf{G} \mathbf{b} = \lambda_b \mathbf{b} \quad (\text{A20})$$

with the resultant secular equation

$$\det(\mathbf{F} \mathbf{G} - \lambda \mathbf{I}) = 0. \quad (\text{A21})$$

From Eq. (A16) we see that the coefficient vector satisfies the normalization condition

$$\mathbf{b}^T \mathbf{G} \mathbf{b} = 1. \quad (\text{A22})$$

B Gramian Determinants

The Gramian determinant[19] is defined as:

$$\Gamma \equiv \det(\gamma_{ij}), \quad (\text{B1})$$

where $\gamma_{ij} = \mathbf{r}_i \cdot \mathbf{r}_j / r_i r_j$, the angle cosines between the particle radii $\mathbf{r}_i$ and $\mathbf{r}_j$, represents the elements of an $N \times N$ matrix. A related quantity used in the main text is the principle minor of the Gramian, $\Gamma^{(\alpha)}$, defined as the determinant of the γ_{ij} matrix with the α^{th} row and column removed.

A most challenging part of calculating the large- D minimum and the $\mathbf{F}$ matrix elements in the systems discussed in this paper is handling the Gramian determinants and their derivatives. What makes these calculations feasible is the very high symmetry of the infinite-dimension, symmetric minimum. In this appendix we discuss the Gramian determinant and show how to obtain its derivatives at the infinite-dimension, symmetric minimum. We will now demonstrate how this is done and summarize the results.

In general $\gamma_{ii} = 1$ and at the infinite- D symmetric minimum all of the remaining direction cosines are equal, $\gamma_{ij} = \bar{\gamma}_\infty$. Hence, the Gramian determinant is an $(N \times N)$ matrix of the form

$$\Gamma|_\infty = \det \begin{pmatrix} 1 & \bar{\gamma}_\infty & \cdots & \bar{\gamma}_\infty \\ \bar{\gamma}_\infty & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \bar{\gamma}_\infty \\ \bar{\gamma}_\infty & \cdots & \bar{\gamma}_\infty & 1 \end{pmatrix}, \quad (\text{B2})$$

in Appendices B and D of Ref. [2] we show that (Eq. (D4))

$$\Gamma|_\infty = [1 + (N-1)\bar{\gamma}_\infty](1 - \bar{\gamma}_\infty)^{N-1}. \quad (\text{B3})$$

The principle minor evaluated at the infinite- D symmetric minimum $\Gamma^{(\alpha)}|_\infty$ is simply related to the corresponding Gramian determinant (B3) by $N \rightarrow N-1$.

To calculate the infinite- D symmetric minimum of the Gramian derivatives, we expand Γ in terms of its cofactors. The cofactor, denoted by C_{ij} , of the element γ_{ij} in Γ is $(-1)^{i+j}$ multiplied by the determinant of the matrix obtained by deleting the i^{th} row and j^{th} column of Γ . We may then write Γ as

$$\Gamma = \sum_{j=1}^N \gamma_{ij} C_{ij}. \quad (\text{B4})$$

Then the partial derivative of Γ in terms of the cofactor is

$$\frac{\partial \Gamma}{\partial \gamma_{ij}} = 2C_{ij}. \quad (\text{B5})$$

From this equation, the partial derivative of Γ evaluated at the infinite- D symmetric minimum is

$$\left. \frac{\partial \Gamma}{\partial \gamma_{ij}} \right|_\infty = -2C_\infty^{(N-1)}, \quad (\text{B6})$$

where we have defined the following determinant of an $(N-1) \times (N-1)$ matrix:

$$C_{\infty}^{(N-1)} = \det \begin{pmatrix} \bar{\gamma}_{\infty} & \bar{\gamma}_{\infty} & \bar{\gamma}_{\infty} & \cdots & \bar{\gamma}_{\infty} \\ \bar{\gamma}_{\infty} & 1 & \bar{\gamma}_{\infty} & & \vdots \\ \bar{\gamma}_{\infty} & \bar{\gamma}_{\infty} & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \bar{\gamma}_{\infty} \\ \bar{\gamma}_{\infty} & \cdots & \cdots & \bar{\gamma}_{\infty} & 1 \end{pmatrix}, \quad (\text{B7})$$

and the superscript $(N-1)$ simply indicates the size of the matrix. From this matrix one can show that the following recursion relation holds:

$$C_{\infty}^{(N-1)} = \bar{\gamma}_{\infty} \left(\Gamma \Big|_{\infty}^{(N-2)} - (N-2)C_{\infty}^{(N-2)} \right), \quad (\text{B8})$$

or equivalently

$$C_{\infty}^{(N)} = \bar{\gamma}_{\infty} \left(\Gamma \Big|_{\infty}^{(N-1)} - (N-1)C_{\infty}^{(N-1)} \right), \quad (\text{B9})$$

where the (N) superscript in the notation $\Gamma \Big|_{\infty}^{(N)}$ again refers to the size of the matrix $\Gamma \Big|_{\infty}$. From the recursion relation (B9), one can easily prove by induction the conjecture that $C_{\infty}^{(N)} = \bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-2}$ and so,

$$\frac{\partial \Gamma}{\partial \gamma_{ij}} \Big|_{\infty} = -2\bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-2}. \quad (\text{B10})$$

The derivative of the principle minor evaluated at the infinite- D symmetric minimum is simply related to the corresponding Gramian determinant derivative (B10) by $N \rightarrow N-1$.

To summarize the above results, the following expressions are needed when calculating the minimum of the effective potential (26 and 27):

$$\begin{aligned} \frac{\partial \Gamma}{\partial \gamma_{ij}} \Big|_{\infty} &= -2\bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-2} & \frac{\partial \Gamma^{(\alpha)}}{\partial \gamma_{ij}} \Big|_{\infty} &= -2\bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-3} \\ \Gamma \Big|_{\infty} &= [1 + (N-1)\bar{\gamma}_{\infty}](1 - \bar{\gamma}_{\infty})^{N-1} & \Gamma^{(\alpha)} \Big|_{\infty} &= [1 + (N-2)\bar{\gamma}_{\infty}](1 - \bar{\gamma}_{\infty})^{N-2}. \end{aligned} \quad (\text{B11})$$

And when evaluating the $\mathbf{F}$ matrix elements at the infinite- D symmetric minimum, the following six second-order derivatives of the Gramian determinants are needed:

$$\begin{aligned} \frac{\partial^2 \Gamma}{\partial \gamma_{ij} \partial \gamma_{kl}} \Big|_{\infty} &= 0 & \frac{\partial^2 \Gamma}{\partial \gamma_{ij}^2} \Big|_{\infty} &= -2(1 - \bar{\gamma}_{\infty})^{N-3}(1 + (N-3)\bar{\gamma}_{\infty}) & \frac{\partial^2 \Gamma}{\partial \gamma_{ij} \partial \gamma_{jk}} \Big|_{\infty} &= 2\bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-3} \\ \frac{\partial^2 \Gamma^{(\alpha)}}{\partial \gamma_{ij} \partial \gamma_{kl}} \Big|_{\infty} &= 0 & \frac{\partial^2 \Gamma^{(\alpha)}}{\partial \gamma_{ij}^2} \Big|_{\infty} &= -2(1 - \bar{\gamma}_{\infty})^{N-4}(1 + (N-4)\bar{\gamma}_{\infty}) & \frac{\partial^2 \Gamma^{(\alpha)}}{\partial \gamma_{ij} \partial \gamma_{jk}} \Big|_{\infty} &= 2\bar{\gamma}_{\infty}(1 - \bar{\gamma}_{\infty})^{N-4}. \end{aligned} \quad (\text{B12})$$

C The Symmetric Group and the Theory of Group Characters

A group of transformations is a set of transformations which satisfy the composition law $ab = c$ where a and b are any two elements of the group and c also belongs to the group. A group also contains the identity element I such that $aI = Ia = a$ and for every element in the group its inverse is also to be found in the group.

The symmetric group S_N is the group of all permutations of N objects and, as such, has $N!$ elements. A permutation may be written as

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 3 & 1 & 5 & 4 & 7 & 6 & 8 \end{pmatrix} \quad (\text{C1})$$

This denotes the following transformation

Object before transformation	$\longrightarrow$	is transformed to object
1	$\longrightarrow$	2
2	$\longrightarrow$	3
3	$\longrightarrow$	1
4	$\longrightarrow$	5
5	$\longrightarrow$	4
6	$\longrightarrow$	7
7	$\longrightarrow$	6
8	$\longrightarrow$	8

A cycle is a particular kind of permutation where the object labels are permuted into each other cyclically. For example the cycle (abc) means that object a is transformed into object b , object b is transformed into object c and object c is transformed into object a . The cycle (abc) is termed a 3-cycle since it cycles three objects. Like wise (3479) is a 4-cycle and (5) is a 1-cycle, the latter transforms object five into itself. All $N!$ permutations of the group S_N may be decomposed into cycles. For example, the permutation of Eq. (C1) may be decomposed into cycles as

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 3 & 1 & 5 & 4 & 7 & 6 & 8 \end{pmatrix} = (123)(45)(67)(8) \quad (\text{C2})$$

This consists of one 1-cycle, two 2-cycles and one 3-cycle. We can denote the cycle structure of a permutation by the symbol $(1^{\nu_1}, 2^{\nu_2}, 3^{\nu_3}, \dots, N^{\nu_N})$, [15] where the notation j^{ν_j} means a cycle of length j and ν_j equals the number of cycles of length j in that permutation. In the case of the permutation of Eq. (C2), its cycle structure is $(1^1, 2^2, 3^1)$.

A matrix representation of a group is a set of nonsingular matrices, including the unit matrix, which has the same composition law as the elements of the group. The character of an element of an matrix representation of a group is the trace of the matrix. The character, as the trace of a matrix, is invariant under a similarity transformation and thus elements of equivalent representations have the same character. The set of all the distinct characters of the elements of an irreducible representation of the group uniquely specify the irreducible representation up to an equivalence transformation. The characters of irreducible representations of a group are termed *simple* characters. All elements of a group which are related by a similarity transformation are said to belong to the same class. The character of the elements of a group

belonging to a particular class all have the same character. Thus there are as many distinct characters for a group as there are classes.

For the group S_N all elements with the same cycle structure belong to the same class and so all elements of a matrix representation of S_N with the same cycle structure have the same character.

A reducible matrix representation of a group may be brought to block diagonal form by a similarity transformation, where the individual blocks are irreducible matrix representations of the same group with lower dimensionality. Thus the characters, $\chi(R)$, of a reducible group are the sums of the characters of the irreducible matrix representations into which it can be decomposed, i.e.

$$\chi(R) = \sum_p \chi_p(R), \quad (\text{C3})$$

where R denotes the element of the group, p labels all of the irreducible blocks into which the reducible matrix representation of the group may be decomposed and $\chi_p(R)$ is the character of the irreducible representation of the block labelled by p . Now in a particular reducible representation a given irreducible representation may be repeated along the diagonal a_α times and so Eq. (C3) may be rewritten as

$$\chi(R) = \sum_\alpha a_\alpha \chi_\alpha(R). \quad (\text{C4})$$

The decomposition of $\chi(R)$ of Eq. (C4) into simple characters $\chi_\alpha(R)$ is unique, i.e. there is not another decomposition of the form

$$\chi(R) = \sum_\alpha b_\alpha \chi_\alpha(R), \quad (\text{C5})$$

where at least one of the b_α is different from the corresponding a_α . This follows from the fact that quite generally

$$a_\alpha = \frac{1}{h} \sum_R \chi_\alpha^*(R) \chi(R) \quad (\text{C6})$$

and that the simple characters satisfy the orthogonality condition

$$\sum_R \chi_\alpha^*(R) \chi_\beta(R) = h \delta_{\alpha\beta}, \quad (\text{C7})$$

where h is the number of elements in the group. For Eqs. (C5), (C6) and (C7) to be consistent we must have

$$b_\alpha = a_\alpha. \quad (\text{C8})$$

The irreducible matrix representations of S_N may be labelled by a Young diagram (= Young pattern = Young shape). A Young diagram is a set of N adjacent squares such that the row below a given row of squares is equal to or shorter in length. The set of all Young diagrams that can be formed from N boxes of all possible irreducible matrix representations of S_N .

A given Young diagram may be denoted by a partition. A partition, $[\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_N]$ is a series of N numbers λ_i such that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_N$ such that $\lambda_1 + \lambda_2 + \lambda_3 + \dots + \lambda_N = N$. The number λ_i is the number of boxes in row i of the Young diagram. Thus the set of all possible partitions of length

N labels all of the possible irreducible matrix representations of S_N and so the irreducible representation labels α and β in Eqs. (C4), (C5), (C6), (C7) and (C8) above, for the group S_N may be taken to run over the set of all possible partitions.

A Young diagram can have up to N rows. For an N -row (one-column) Young diagram all of the λ_i s are non zero. However only one Young diagram will have N row; all of the rest will have less than N rows. Thus in all but one case the last few λ_i s will be zero. It is standard practice to drop the zeros and use a partition with less than N numbers. Thus the partition $[3, 0, 0]$ labelling an irreducible representation of S_3 is usually abbreviated to just $[3]$.

D The Inequivalence of Subspaces α and α' under S_N when $\alpha \neq \alpha'$ – A Useful Equation.

Here we wish to note something that proves useful in Sec. 6.2 above. Consider the quantity

$$W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T D_{S_{\mathbf{X}'}}^{\alpha'}(R), \quad (\text{D1})$$

where $\mathbf{X}'$ is $\bar{\mathbf{r}}'$ or $\bar{\mathbf{y}}'$, α and α' denote the partitions $[N]$, $[N-1, 1]$ or $[N-2, 2]$, and Υ is an unspecified matrix. The matrix $W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T$ effects a similarity mapping from the irreducible α' and $S_{\mathbf{X}'}$ space onto an irreducible space denoted by α . Thus one can construct an irreducible matrix representation $D_c^\alpha(R)$ belonging to the set of equivalent representations denoted by α such that

$$D_c^\alpha(R) W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T = W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T D_{S_{\mathbf{X}'}}^{\alpha'}(R) \quad (\text{D2})$$

for all elements R of the group. According to Schur's Lemma however, any such similarity mapping, $W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T$, that satisfies Eq. (D2) is identically zero unless $\alpha = \alpha'$, i.e.

$$W_{\mathbf{X}'}^\alpha \Upsilon [W_{\mathbf{X}'}^{\alpha'}]^T = 0, \quad \forall \quad \alpha \neq \alpha'. \quad (\text{D3})$$

Equation (D3) is a statement of the inequivalence of the irreducible spaces labelled by α and α' under S_N when $\alpha \neq \alpha'$. In particular we have

$$W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^{\alpha'}]^T = 0, \quad \forall \quad \alpha \neq \alpha'. \quad (\text{D4})$$

Schur's Lemma also implies $D_c^{\alpha'}(R) \neq D_{S_{\mathbf{X}'}}^{\alpha'}(R)$ unless $W_{\mathbf{X}'}^{\alpha'} [W_{\mathbf{X}'}^{\alpha'}]^T \propto I_{\alpha'}$, where $I_{\alpha'}$ is the unit matrix.

E Proof of the Orthogonality of W .

We shall show that condition 2 of Sec 6.2 above implies that W is an orthogonal matrix. This can be seen as follows. The displacement vector $\bar{\mathbf{y}}'$ of Eq. (38) transforms under an orthogonal, though reducible,

representation of S_N since $\bar{\mathbf{y}}'^T \bar{\mathbf{y}}'$ is invariant under S_N . Writing this orthogonal representation as

$$D_{\bar{\mathbf{y}}'}(R) = \left(\begin{array}{c|c} D_{\bar{\mathbf{r}}'}(R) & \mathbf{0} \\ \hline \mathbf{0} & D_{\bar{\mathbf{r}}'}(R) \end{array} \right), \quad (\text{E1})$$

Eq. (100) implies that the representation under which the displacement coordinate vector $\mathbf{S}$ transforms is

$$D_S = W D_{\bar{\mathbf{y}}'}(R) W^{-1}. \quad (\text{E2})$$

Since $D_{\bar{\mathbf{y}}'}(R)$ is an orthogonal matrix representations of S_N then from Eq. (E2) we have

$$D_{\bar{\mathbf{y}}'}^T(R) D_{\bar{\mathbf{y}}'}(R) = W^T D_S^T(R) [W^{-1}]^T W^{-1} D_S(R) W = 1. \quad (\text{E3})$$

From Eq. (E3) we can write

$$D_S^T(R) [W W^T]^{-1} D_S(R) = [W W^T]^{-1} = D_S^T(R) D_S(R) [W W^T]^{-1}, \quad (\text{E4})$$

where we have used the orthogonality of $D_S(R)$ in the last step. Left multiplying by $[D_S^T]^{-1}(R)$ we obtain

$$[W W^T]^{-1} D_S(R) = D_S(R) [W W^T]^{-1}, \quad (\text{E5})$$

from which we derive

$$W W^T D_S(R) = D_S(R) W W^T. \quad (\text{E6})$$

We notice that

$$W W^T = I \quad (\text{E7})$$

satisfies Eq. (E6). We will now show that it is the only solution. We note that $D_S(R)$ is a block diagonal matrix of irreducible representations of the form

$$\begin{aligned} D_S(R) &= P \left(\begin{array}{c|c} D_{S_{\bar{\mathbf{r}}'}}(R) & \mathbf{0} \\ \hline \mathbf{0} & D_{S_{\bar{\mathbf{r}}'}}(R) \end{array} \right) P^T \\ &= \left(\begin{array}{cc|cc|c} D^{[N]}(R) & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & D^{[N]}(R) & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & D^{[N-1, 1]}(R) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & D^{[N-1, 1]}(R) & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & D^{[N-2, 2]}(R) \end{array} \right), \quad (\text{E8}) \end{aligned}$$

where we have also used condition 1 from Sec. 6.2 above in the last step. Now from Eq. (99)

$$\begin{aligned} W W^T &= \\ &\left(\begin{array}{ccccc} W_{\bar{\mathbf{r}}'}^{[N]} [W_{\bar{\mathbf{r}}'}^{[N]}]^T & \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N]} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N]} [W_{\bar{\mathbf{r}}'}^{[N]}]^T & \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N]} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T & W_{\bar{\mathbf{r}}'}^{[N]} [W_{\bar{\mathbf{r}}'}^{[N-2, 2]}]^T \\ W_{\bar{\mathbf{r}}'}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^{[N]}]^T & \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^{[N]}]^T & \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T & W_{\bar{\mathbf{r}}'}^{[N-1, 1]} [W_{\bar{\mathbf{r}}'}^{[N-2, 2]}]^T \\ \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N-2, 2]} [W_{\bar{\mathbf{r}}'}^{[N]}]^T & \mathbf{0} & W_{\bar{\mathbf{r}}'}^{[N-2, 2]} [W_{\bar{\mathbf{r}}'}^{[N-1, 1]}]^T & W_{\bar{\mathbf{r}}'}^{[N-2, 2]} [W_{\bar{\mathbf{r}}'}^{[N-2, 2]}]^T \end{array} \right) \quad (\text{E9}) \end{aligned}$$

Substituting Eqs. (E8) and (E9) in Eq. (E6) we obtain

$$W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^{\alpha'}]^T D^{\alpha'}(R) = D^\alpha(R) W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^{\alpha'}]^T, \quad (\text{E10})$$

where $\mathbf{X}'$ is $\bar{\mathbf{r}}'$ or $\bar{\mathbf{r}}'$, and α and α' are the partitions $[N]$, $[N-1, 1]$ or $[N-2, 2]$. As we note in Appendix D, Eq. (D4), $W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^{\alpha'}]^T = 0$ for $\alpha \neq \alpha'$, and so Eq. (E10) is automatically satisfied when $\alpha \neq \alpha'$. It does not give us a new condition on the $W_{\mathbf{X}'}^\alpha$. When $\alpha = \alpha'$ however, we have

$$W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^\alpha]^T D^\alpha(R) = D^\alpha(R) W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^\alpha]^T. \quad (\text{E11})$$

The second part of Schur's Lemma[20] then requires that

$$W_{\mathbf{X}'}^\alpha [W_{\mathbf{X}'}^\alpha]^T = I_\alpha, \quad (\text{E12})$$

where I_α is the unit matrix.

Thus Eq. (E12) is the essential equation to satisfy.

When Eq. (E12) is satisfied, Eq. (E12), together with the automatically satisfied Eq. (D4), ensures that Eq. (E7) is satisfied. Multiplying Eq. (E7) on the right by W^T and on the left by $[W^T]^{-1}$ we obtain $W^T W = I$ and so W is an orthogonal transformation satisfying

$$W W^T = W^T W = I. \quad (\text{E13})$$

F Calculation of $[\sigma_{[\mathbf{N}]}^{\mathbf{G}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$ and $[\sigma_{[\mathbf{N}]}^{\mathbf{FG}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$, the reduced $\mathbf{G}$ and $\mathbf{FG}$ matrix elements, in the symmetry-coordinate basis.

In this appendix we use the $W_{\mathbf{X}'}^{[N]}$ matrices ($\mathbf{X}' = \bar{\mathbf{r}}'$ or $\bar{\mathbf{r}}'$) to calculate the reduced $\mathbf{G}$ and $\mathbf{FG}$ matrix elements, $[\sigma_{[\mathbf{N}]}^{\mathbf{G}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$ and $[\sigma_{[\mathbf{N}]}^{\mathbf{FG}}]_{\mathbf{x}'_1, \mathbf{x}'_2}$, using Eq. (120):

$$[\sigma_{[\mathbf{N}]}^{\mathbf{Q}}]_{\mathbf{x}'_1, \mathbf{x}'_2} = (W_{\mathbf{x}'_1}^{[N]})_\xi \mathbf{Q}_{\mathbf{x}'_1 \mathbf{x}'_2} [(W_{\mathbf{x}'_2}^{[N]})_\xi]^T, \quad (\text{F1})$$

where $\mathbf{Q} = \mathbf{G}$ or $\mathbf{FG}$ and ξ is a row label.

Using Eq. (188)

$$[W_{\bar{\mathbf{r}}'}^{[N]}]_i = \frac{1}{\sqrt{N}} [\mathbf{1}_{\bar{\mathbf{r}}'}]_i \quad (\text{F2})$$

and Eq. (197)

$$[W_{\bar{\mathbf{r}}'}^{[N]}]_{ij} = \sqrt{\frac{2}{N(N-1)}} [\mathbf{1}_{\bar{\mathbf{r}}'}]_{ij} \quad (\text{F3})$$

with Eq. (66)

$$\mathbf{G} = \begin{pmatrix} \tilde{a}' \mathbf{I}_N & \mathbf{0} \\ \mathbf{0} & \tilde{g}' \mathbf{I}_M + \tilde{h}' \mathbf{R}^T \mathbf{R} \end{pmatrix}, \quad (\text{F4})$$

where $\mathbf{I}_N$, $\mathbf{I}_M$, $\mathbf{R}$ are defined in Sect. 5.3 after Eq. (56) and $\tilde{g}'$ and $\tilde{h}'$ are defined by Eq. (65), we find:

$$\begin{aligned}
[\sigma_{[N]}^G]_{\bar{r}', \bar{r}'} &= \sum_{i,j=1}^N [W_{\bar{r}'}^{[N]}]_i [\mathbf{G}_{\bar{r}' \bar{r}'}]_{ij} [(W_{\bar{r}'}^{[N]})^T]_j \\
&= \frac{\tilde{a}'}{N} \sum_{i,j=1}^N [\mathbf{1}_{\bar{r}'}]_i \delta_{ij} [\mathbf{1}_{\bar{r}'}]_j \\
&= \frac{\tilde{a}'}{N} \sum_i 1 \\
&= \tilde{a}',
\end{aligned} \tag{F5}$$

$$\begin{aligned}
[\sigma_{[N]}^G]_{\bar{r}', \bar{\gamma}'} &= \sum_{i=1}^N \sum_{k=2}^N \sum_{j=1}^{k-1} [W_{\bar{r}'}^{[N]}]_i [\mathbf{G}_{\bar{r}' \bar{\gamma}'}]_{i,j,k} [(W_{\bar{\gamma}'}^{[N]})^T]_{jk} \\
&= 0,
\end{aligned} \tag{F6}$$

$$\begin{aligned}
[\sigma_{[N]}^G]_{\bar{\gamma}', \bar{r}'} &= \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{k=1}^N [W_{\bar{\gamma}'}^{[N]}]_{ij} [\mathbf{G}_{\bar{\gamma}' \bar{r}'}]_{ij,k} [(W_{\bar{r}'}^{[N]})^T]_k \\
&= 0
\end{aligned} \tag{F7}$$

and

$$\begin{aligned}
[\sigma_{[N]}^G]_{\bar{\gamma}', \bar{\gamma}'} &= \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{l=2}^N \sum_{k=1}^{l-1} [W_{\bar{\gamma}'}^{[N]}]_{ij} [\mathbf{G}_{\bar{\gamma}' \bar{\gamma}'}]_{ij,kl} [(W_{\bar{\gamma}'}^{[N]})^T]_{kl} \\
&= \frac{2}{N(N-1)} \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{l=2}^N \sum_{k=1}^{l-1} [\mathbf{1}_{\bar{\gamma}'}]_{ij} (\tilde{g}'(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) + \tilde{h}'(\delta_{ik} + \delta_{il} + \delta_{jk} + \delta_{jl})) [\mathbf{1}_{\bar{\gamma}'}]_{kl} \\
&= \frac{2}{4N(N-1)} \left(\sum_{i,j,k,l=1}^N [\tilde{g}'(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) + \tilde{h}'(\delta_{ik} + \delta_{il} + \delta_{jk} + \delta_{jl})] \right. \\
&\quad - \sum_{i,j,k=1}^N [\tilde{g}'(\delta_{ik}\delta_{jk} + \delta_{ik}\delta_{jk}) + \tilde{h}'(\delta_{ik} + \delta_{ik} + \delta_{jk} + \delta_{jk})] \\
&\quad - \sum_{i,k,l=1}^N [\tilde{g}'(\delta_{ik}\delta_{il} + \delta_{il}\delta_{ik}) + \tilde{h}'(\delta_{ik} + \delta_{il} + \delta_{ik} + \delta_{il})] \\
&\quad \left. + \sum_{i,k=1}^N [\tilde{g}'(\delta_{ik}\delta_{ik} + \delta_{ik}\delta_{ik}) + \tilde{h}'(\delta_{ik} + \delta_{ik} + \delta_{ik} + \delta_{ik})] \right) \\
&= \frac{1}{2N(N-1)} [(2\tilde{g}'N^2 - 4\tilde{h}'N^3) - 2(2\tilde{g}'N + 4\tilde{h}'N^2) + (2\tilde{g}'N + 4\tilde{h}'N)] \\
&= \tilde{g}' + 2\tilde{h}'(N-1).
\end{aligned} \tag{F8}$$

Thus we obtain Eq. (201):

$$\boldsymbol{\sigma}_{[N]}^G = \begin{pmatrix} [\boldsymbol{\sigma}_{[N]}^G]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} = \tilde{a}' & [\boldsymbol{\sigma}_{[N]}^G]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} = 0 \\ [\boldsymbol{\sigma}_{[N]}^G]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} = 0 & [\boldsymbol{\sigma}_{[N]}^G]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} = (\tilde{g}' + 2(N-1)\tilde{h}') \end{pmatrix}. \quad (\text{F9})$$

Similarly letting $\mathbf{Q} = \mathbf{F}\mathbf{G}$ and using Eq. (62):

$$\mathbf{F}\mathbf{G} = \begin{pmatrix} \tilde{a}\mathbf{I}_N + \tilde{b}\mathbf{J}_N & \tilde{e}\mathbf{R} + \tilde{f}\mathbf{J}_{NM} \\ \tilde{c}\mathbf{R}^T + \tilde{d}\mathbf{J}_{MN} & \tilde{g}\mathbf{I}_M + \tilde{h}\mathbf{R}^T\mathbf{R} + \tilde{l}\mathbf{J}_M \end{pmatrix}, \quad (\text{F10})$$

we can derive:

$$\begin{aligned} [\boldsymbol{\sigma}_{[N]}^{FG}]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} &= \sum_{i,j=1}^N [W_{\bar{\mathbf{r}}'}^{[N]}]_i [\mathbf{F}\mathbf{G}_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'}]_{ij} [(W_{\bar{\mathbf{r}}'}^{[N]})^T]_j \\ &= \frac{1}{N} \sum_{i,j=1}^N [\mathbf{1}_{\bar{\mathbf{r}}'}]_i (\tilde{a}\delta_{ij} + \tilde{b}\mathbf{1}_{ij}) [\mathbf{1}_{\bar{\mathbf{r}}'}]_j \\ &= \frac{1}{N} [N\tilde{a} + \tilde{b}N^2] \\ &= \tilde{a} + \tilde{b}N, \end{aligned} \quad (\text{F11})$$

$$\begin{aligned} [\boldsymbol{\sigma}_{[N]}^{FG}]_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'} &= \sum_{i=1}^N \sum_{k=2}^N \sum_{j=1}^{k-1} [W_{\bar{\mathbf{r}}'}^{[N]}]_i [\mathbf{F}\mathbf{G}_{\bar{\mathbf{r}}', \bar{\mathbf{r}}'}]_{i,jk} [(W_{\bar{\mathbf{r}}'}^{[N]})^T]_{jk} \\ &= \frac{1}{N} \sqrt{\frac{2}{N-1}} \sum_{i=1}^N \sum_{k=2}^N \sum_{j=1}^{k-1} [\mathbf{1}_{\bar{\mathbf{r}}'}]_i \left(\tilde{e}(\delta_{ij} + \delta_{ik}) + \tilde{f} \right) [\mathbf{1}_{\bar{\mathbf{r}}'}]_{jk} \\ &= \frac{1}{N\sqrt{2(N-1)}} \left(\sum_{i,j,k=1}^N \tilde{e}(\delta_{ij} + \delta_{ik}) + \tilde{f} - \sum_{i,j=1}^N \tilde{e}(\delta_{ij} + \delta_{ij}) + \tilde{f} \right) \\ &= \frac{1}{N\sqrt{2(N-1)}} (2\tilde{e}N^2 + \tilde{f}N^3 - 2\tilde{e}N - \tilde{f}N^2) \\ &= \sqrt{2(N-1)} \left(\tilde{e} + \frac{N}{2}\tilde{f} \right), \end{aligned} \quad (\text{F12})$$

$$\begin{aligned}
[\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{r}'} &= \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{k=1}^N [W_{\bar{\gamma}'}^{[N]}]_{ij} [\mathbf{F}G_{\bar{\gamma}' \bar{r}'}]_{ij, k} [(W_{\bar{r}'}^{[N]})^T]_k \\
&= \frac{1}{N} \sqrt{\frac{2}{N-1}} \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{k=1}^N [\mathbf{1}_{\bar{\gamma}'}]_{ij} [\tilde{c}(\delta_{ki} + \delta_{kj}) + \tilde{d}] [\mathbf{1}_{\bar{r}'}]_k \\
&= \frac{1}{N\sqrt{2(N-1)}} \left(\sum_{i,j,k=1}^N [\tilde{c}(\delta_{ki} + \delta_{kj}) + \tilde{d}] - \sum_{ik=1}^N [\tilde{c}(\delta_{ki} + \delta_{ki}) + \tilde{d}] \right) \\
&= \frac{1}{N\sqrt{2(N-1)}} (2\tilde{c}N^2 + \tilde{d}N^3 - 2\tilde{c}N - \tilde{d}N^2) \\
&= \sqrt{2(N-1)} \left(\tilde{c} + \frac{N}{2} \tilde{d} \right) \tag{F13}
\end{aligned}$$

and

$$\begin{aligned}
[\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} &= \sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{l=2}^N \sum_{k=1}^{l-1} [W_{\bar{\gamma}'}^{[N]}]_{ij} [\mathbf{F}G_{\bar{\gamma}' \bar{\gamma}'}]_{ij, kl} [(W_{\bar{\gamma}'}^{[N]})^T]_{kl} \\
&= \frac{2}{N(N-1)} \left(\sum_{j=2}^N \sum_{i=1}^{j-1} \sum_{l=2}^N \sum_{k=1}^{l-1} [\mathbf{1}_{\bar{\gamma}'}]_{ij} [\tilde{g}(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) + \tilde{h}(\delta_{ik} + \delta_{il} + \delta_{jk} + \delta_{jl}) + \tilde{i}] [\mathbf{1}_{\bar{\gamma}'}]_{kl} \right) \\
&= \frac{1}{2N(N-1)} \left(\sum_{i,j,k,l=1}^N [\tilde{g}(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) + \tilde{h}(\delta_{ik} + \delta_{il} + \delta_{jk} + \delta_{jl}) + \tilde{i}] \right. \\
&\quad - \sum_{i,j,k=1}^N (\tilde{g}(\delta_{ik}\delta_{jk} + \delta_{ik}\delta_{jk}) + \tilde{h}(\delta_{ik} + \delta_{ik} + \delta_{jk} + \delta_{jk}) + \tilde{i}) \\
&\quad - \sum_{i,k,l=1}^N (\tilde{g}(\delta_{ik}\delta_{il} + \delta_{il}\delta_{ik}) + \tilde{h}(\delta_{ik} + \delta_{il} + \delta_{ik} + \delta_{il}) + \tilde{i}) \\
&\quad \left. + \sum_{i,k=1}^N (\tilde{g}(\delta_{ik}\delta_{ik} + \delta_{ik}\delta_{ik}) + \tilde{h}(\delta_{ik} + \delta_{ik} + \delta_{ik} + \delta_{ik}) + \tilde{i}) \right) \\
&= \frac{1}{2N(N-1)} (2N(N-1)\tilde{g} + 4N(N^2 - 2N + 1)\tilde{h} + N^2(N^2 - 2N + 1)\tilde{i}) \\
&= \tilde{g} + 2(N-1)\tilde{h} + \frac{N(N-1)}{2} \tilde{i}. \tag{F14}
\end{aligned}$$

Thus we obtain Eq. (202):

$$\sigma_{[N]}^{FG} = \begin{pmatrix} [\sigma_{[N]}^{FG}]_{\bar{r}', \bar{r}'} = (\tilde{a} + \tilde{b}N) & [\sigma_{[N]}^{FG}]_{\bar{r}', \bar{\gamma}'} = \sqrt{2(N-1)} \left(\tilde{e} + \frac{N}{2} \tilde{f} \right) \\ [\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{r}'} = \sqrt{2(N-1)} \left(\tilde{c} + \frac{N}{2} \tilde{d} \right) & [\sigma_{[N]}^{FG}]_{\bar{\gamma}', \bar{\gamma}'} = \left(\tilde{g} + 2(N-1)\tilde{h} + \frac{N(N-1)}{2} \tilde{i} \right) \end{pmatrix}. \tag{F15}$$

References

- [1]
- [2] B.A. McKinney, M. Dunn, D.K. Watson, and J.G. Loeser, *Ann. Phys.* **310**, 56 (2003).
- [3] D.R. Herschbach, J. Avery, and O. Goscinski, Eds., *Dimensional Scaling in Chemical Physics*. Kluwer, Dordrecht, 1992.
- [4] B.A. McKinney, M. Dunn, D.K. Watson, *Phys. Rev. A* **69**, 053611 (2004).
- [5] D. Blume and C.H. Greene, *Phys. Rev. A* **63**, 63061 (2001).
- [6] M. Dunn, T.C. Germann, D.Z. Goodson, C.A. Traynor, J.D. Morgan III, D.K. Watson, and D.R. Herschbach, *J. Chem. Phys.* **101**, 5987 (1994).
- [7] S.L. Cornish, N.R. Claussen, J.L. Roberts, E.A. Cornell, and C.E. Wieman, *Phys. Rev. Lett.* **85**, 1795 (2000).
- [8] J.G. Loeser, *J. Chem. Phys.* **86**, 5635 (1987).
- [9] W.B. Laing, M. Dunn, J.G. Loeser and D.K. Watson, submitted to *Phys. Rev. Lett.*
- [10] M. Dunn, D.K. Watson, and J.G. Loeser, to be submitted *Ann. Phys.*
- [11] J. Avery, D.Z. Goodson, D.R. Herschbach, *Theor. Chim. Acta* **81**, 1 (1991).
- [12] A. Chatterjee, *J. Phys. A: Math. Gen.* **18**, 735 (1985).
- [13] G. Strang, *Linear Algebra and its Applications, Third ed.*. Harcourt Brace Jovanovich College Publishers, Orlando, FL, 1988.
- [14] E.B. Wilson, Jr., J.C. Decius, P.C. Cross, *Molecular vibrations: The theory of infrared and raman vibrational spectra*. McGraw-Hill, New York, 1955.
- [15] M. Hamermesh, *Group theory and its application to physical problems*. Addison-Wesley, Reading, MA, 1962.
- [16] In Ref. [8], Eq. (53) would read

$$\begin{aligned}
 \overline{E} &= \overline{E}_\infty + \delta \overline{E}_o + O(\delta^2) \\
 &= V_{\text{eff}}(\bar{r}_\infty, \bar{\gamma}_\infty) + \delta \left\{ \sum_{\mu=\{\mathbf{0}^\pm, \mathbf{1}^\pm, \mathbf{2}^\pm\}} (n_\mu + \frac{d_\mu}{2}) \bar{\omega}_\mu + v_o \right\} + O(\delta^2),
 \end{aligned} \tag{F16}$$

where n_μ is the total number of quanta in all the normal modes with the same frequency $\bar{\omega}_\mu$, i.e.

$$n_\mu = \sum_{n_\mu=0}^{\infty} n_\mu d_{\mu, n_\mu}. \tag{F17}$$

- [17] Predrag Cvitanovic, *Group Theory Part I*. Nordita, Copenhagen, 1984, p. 71.
- [18] See for example Ref. [14], Appendix XII, p. 347.
- [19] F.R. Gantmacher, *The theory of matrices, Vol. 1*. Chelsea, New York, 1959.
- [20] See for example Ref. [15], p. 100.